

Epithelial Upregulation of *Toll-like Receptor 2 (TLR2)* Fast-tracks Tissue Repair in ICR Mice Treated with Postbiotic Hydrogel Infused with *Lactobacillus paracasei* 7060



Althea Gay B. Pagurayan¹ , Jimmbeth Zenila P. Fabia^{1,2} , Ma. Joy Theresa T. Agcaoili^{1,2} , James Patrick A. Acang^{1,4} , James Patrick K. Peralta^{1,5} , Alvin Domingo³  and Peter James Icalia-Gann^{1,6,*} 

¹Center for Cellular and Molecular Medical Research, Mariano Marcos State University, Batac 2906, Ilocos Norte, Batac City, Philippines

²Research Directorate, Mariano Marcos State University, Batac 2906, Ilocos Norte, Batac City, Philippines

³Department of Biological Sciences, College of Arts and Sciences, Mariano Marcos State University, Batac 2906, Ilocos Norte, Batac City, Philippines

⁴Department of Computer Science, College of Computing and Information Sciences, Mariano Marcos State University, Batac 2906, Ilocos Norte, Batac City, Philippines

⁵College of Medicine, Mariano Marcos State University, Batac 2906, Ilocos Norte, Batac City, Philippines

⁶Department of Biochemistry and Molecular Biology, College of Medicine, Mariano Marcos State University, Batac 2906, Ilocos Norte, Batac City, Philippines

Abstract:

Introduction: Impaired epithelial wound repair contributes to increased infection risk and delayed tissue recovery. Limitations associated with conventional topical therapies, including antimicrobial resistance, highlight the need for alternative strategies. Postbiotics, defined as bioactive microbial metabolites, have emerged as potential modulators of tissue repair and immune responses.

Methods: A postbiotic hydrogel formulated with metabolites derived from *Lactobacillus paracasei* 7060 was applied to full-thickness dorso-lumbar wounds in a murine model. Wound closure was assessed over a 21-day period. Histological analyses evaluated epithelialization and tissue organization, and expression of immune- and repair-associated genes was quantified by quantitative PCR.

Results: Treatment with the postbiotic hydrogel resulted in accelerated wound closure and earlier epithelial restoration compared with both the commercial hydrogel and untreated control groups. Histological assessment demonstrated improved epidermal continuity and reduced inflammatory cell infiltration in treated wounds. Gene expression analysis revealed a transient upregulation of *Toll-Like Receptor 2 (TLR2)* during the early phase of healing.

Discussion: The observed transient activation of *TLR2* suggests a regulated immune response that may support early host defense and macrophage recruitment without sustained inflammation. Genomic features of *L. paracasei* 7060 are consistent with the production of metabolites involved in immune modulation, tissue repair, angiogenesis, and stress response pathways.

Conclusion: These findings indicate that a *L. paracasei* 7060-derived postbiotic hydrogel supports epithelial wound repair through coordinated immune and regenerative mechanisms.

Keywords: Probiotics, Postbiotics, Tissue repair, Toll-like receptor, Hydrogel, Immunomodulatory biomarkers.

© 2026 The Author(s). Published by Bentham Open.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



Received: October 11, 2025
Revised: December 03, 2025
Accepted: December 11, 2025
Published: March 03, 2026



Send Orders for Reprints to
reprints@benthamscience.net

*Address correspondence to this author at the Center for Cellular and Molecular Medical Research, Mariano Marcos State University, Batac 2906, Ilocos Norte, Batac City, Philippines and Department of Biochemistry and Molecular Biology, College of Medicine, Mariano Marcos State University, Batac 2906, Ilocos Norte, Batac City, Philippines;
E-mail: pjicalia@gmail.com

Cite as: Pagurayan A, Fabia J, Agcaoili M, Acang J, Peralta J, Domingo A, Icalia-Gann P. Epithelial Upregulation of Toll-like Receptor 2 (TLR2) Fast-tracks Tissue Repair in ICR Mice Treated with Postbiotic Hydrogel Infused with *Lactobacillus paracasei* 7060. Open Microbiol J, 2026; 20: e18742858448770. <http://dx.doi.org/10.2174/0118742858448770260210051054>

1. INTRODUCTION

Epithelial tissues serve as the body's first line of defense against environmental stressors, microbial invasion, and mechanical injury, while also supporting vital physiological functions such as absorption, secretion, and immune surveillance [1]. These tissues line the skin and mucosal surfaces, positioning them at the interface between the host and its environment. When epithelial integrity is disrupted, whether in the skin, gastrointestinal tract, or respiratory system, the body initiates a highly coordinated wound healing response involving inflammation, cellular proliferation, angiogenesis, and remodeling of the extracellular matrix (ECM). Precise regulation of each of these stages is essential, as failure to resolve injury appropriately can lead to chronic wounds, excessive scarring, fibrosis, or systemic infection, all of which significantly contribute to morbidity and mortality [2].

Loss of epithelial integrity markedly increases susceptibility to microbial invasion. Open wounds, burns, and damaged mucosal surfaces provide direct portals of entry for opportunistic pathogens, including antibiotic-resistant Gram-negative and Gram-positive bacteria as well as fungal species [3]. Once established, these infections intensify local inflammation, delay tissue regeneration, and can progress to life-threatening conditions such as sepsis if left unchecked. The clinical challenge is further compounded by the increasing prevalence of multidrug-resistant organisms, including *Staphylococcus aureus*, *Escherichia coli*, and *Candida albicans*, which frequently exhibit reduced responsiveness to conventional antimicrobial therapies. This growing resistance crisis underscores the urgent need for alternative therapeutic approaches that can simultaneously enhance host defense mechanisms and limit microbial burden at the wound site [4].

Inflammation plays a critical role in the early phase of wound healing by facilitating pathogen clearance and debris removal. However, when inflammatory responses become excessive or prolonged, they can perpetuate tissue damage rather than promote repair. Elevated concentrations of pro-inflammatory cytokines such as Tumor Necrosis Factor-alpha (TNF- α) and Interleukin-6 (IL-6), together with increased production of Reactive Oxygen Species (ROS), contribute to ECM degradation and interfere with epithelial cell proliferation and

migration [5]. Chronic inflammatory disorders, including diabetic ulcers, inflammatory bowel disease, and chronic obstructive pulmonary disease, illustrate how unresolved epithelial injury leads to sustained tissue dysfunction and impaired healing. Persistent epithelial damage may also progress to fibrosis, a pathological condition characterized by excessive ECM deposition, particularly collagen, driven by the activation of myofibroblasts and regulated by signaling pathways involving Transforming Growth Factor-beta (TGF- β), integrins, and growth factor receptors [6, 7]. Although fibrosis initially serves to stabilize damaged tissue, uncontrolled fibrotic responses can result in disfiguring outcomes such as hypertrophic scars and keloids. Consequently, there is a critical need for therapeutic strategies that support regenerative healing while limiting excessive inflammation and fibrotic remodeling.

In recent years, increasing attention has been directed toward the microbiome and its bioactive derivatives as modulators of epithelial repair. Probiotics, defined as live microorganisms that confer health benefits to the host, have demonstrated promising effects in various wound healing and inflammatory models, largely due to their antimicrobial and immunomodulatory properties. Several studies have shown that *Lactobacillus* and *Bifidobacterium* species can enhance re-epithelialization, stimulate collagen deposition, and regulate cytokine production. For example, *L. rhamnosus* GG and *B. animalis* subsp. *lactis* BB12 was reported to accelerate granulation tissue formation while suppressing pro-inflammatory mediators such as TNF- α and IL-6, thereby promoting macrophage polarization toward an anti-inflammatory M2 phenotype [8, 9]. Topical application of *L. plantarum* protein fractions has been shown to enhance keratinocyte migration and modulate cytokine expression within wound beds [9], while oral supplementation with *L. fermentum*, *L. casei*, and *B. bifidum* reduced oxidative stress in inflammatory conditions [10]. Despite these benefits, the clinical use of live probiotics remains controversial, particularly in immunocompromised individuals or in open wound environments, where bacterial translocation and systemic infection represent potential safety concerns [11].

These limitations have driven growing interest in postbiotics, which are defined by the International Scientific Association for Probiotics and Prebiotics (ISAPP) as "preparations of inanimate microorganisms and/or their

components that confer a health benefit on the host" [12]. Postbiotics encompass a wide range of microbial-derived molecules, including Short-Chain Fatty Acids (SCFAs), exopolysaccharides, bacteriocins, teichoic acids, peptidoglycans, lipoteichoic acids, enzymes, and heat-inactivated bacterial cells [13]. Accumulating evidence indicates that postbiotics can enhance epithelial barrier integrity through the upregulation of tight junction proteins [14], stimulate fibroblast proliferation, keratinocyte migration, and angiogenesis [15], and modulate immune responses by increasing anti-inflammatory cytokines such as IL-10 and TGF- β while suppressing pro-inflammatory mediators including TNF- α and IL-6 [16]. In addition, certain postbiotic components exhibit antimicrobial activity by disrupting pathogenic biofilms and interfering with quorum-sensing mechanisms [17].

Preclinical studies have demonstrated the capacity of postbiotic formulations to accelerate wound healing processes. For instance, supernatants derived from *L. plantarum* have been shown to enhance granulation tissue formation and promote re-epithelialization in cutaneous wound models [18]. Similarly, a lysate of *L. paracasei* CNCM I-1518 improved skin barrier function and reduced inflammatory symptoms in patients with atopic dermatitis [19, 20]. Compared with live probiotics, postbiotics offer several practical advantages, including improved safety profiles for high-risk populations, standardized composition and dosing, enhanced stability and shelf life without the need for refrigeration, and the absence of risks associated with microbial overgrowth or horizontal transfer of antibiotic resistance genes [21-23].

Despite their growing therapeutic potential, investigations into postbiotics for tissue regeneration remain limited, particularly those derived from plant-associated Lactic Acid Bacteria (LAB). Most existing studies focus on strains originating from the human gastrointestinal tract, leaving a substantial gap in knowledge regarding LAB isolated from plant sources. This gap is especially relevant for microbes derived from *Nypa fruticans* (nipa palm), an indigenous species found in Philippine mangrove ecosystems that is recognized for its rich and unique microbial diversity. In this study, we investigate the wound healing potential of a postbiotic hydrogel derived from *Lactobacillus paracasei* 7060, a strain isolated from nipa palm. Using phenotypic wound assessments, histological evaluation, and analysis of key inflammation- and regeneration-related genes (TLR2, TNF- α , and TGF- β), we aim to assess the capacity of this plant-derived postbiotic to favorably modulate the wound microenvironment. By integrating indigenous microbial resources with translational biomedical research, this work provides novel insights into the role of postbiotics in epithelial regeneration and immune modulation.

2. MATERIALS AND METHODS

2.1. Research Design

A Completely Randomized Design (CRD) was employed with five intervention groups and six biological replicates

each. Blocking was based on tissue harvest time: half of each group was sacrificed on day 3 post-wounding, and the remaining on day 21. All mice (except the basal control) underwent wound excision under aseptic conditions. Thirty male ICR mice were randomly assigned to the following groups:

Basal Control - No wound or treatment

Experimental - Wound treated with *postbiotic*-formulated hydrogel

Negative Control 1 - Wound treated with hydrogel base only

Negative Control 2 - Wound without any treatment

Positive Control - Wound treated with DuoDERM hydrogel + povidone-iodine

2.2. Preparation of *Postbiotic*-formulated Hydrogel

Powdered nipa fronds underwent soda treatment followed by sequential NaOH washing, filtration, bleaching, and drying to yield α -cellulose. Carboxymethylcellulose (CMC) was synthesized from purified cellulose. A chilled alkaline/urea solution (7% NaOH, 12% urea, 81% H₂O) was pre-cooled to -12.6°C. CMC (1 g) was added and stirred (1500 rpm, 10 min), followed by centrifugation (8000 rpm, 20 min), neutralization with 10% H₂SO₄, and autoclaving. *L. paracasei* 7060 was cultured in MRS broth (35-37°C, 48 hours). Aseptically, 50 mL CMC solution was combined with sterilized carbomer while stirring until homogeneous. Gelation was initiated with triethanolamine, followed by the addition of 50 mL probiotic-inoculated MRS broth. The final hydrogel was transferred into sterile petri dishes and stored at room temperature.

2.3. *In Vitro* Antibacterial Assay of *Postbiotic*-formulated Hydrogel

Mueller-Hinton Agar (MHA) was prepared, sterilized, and poured into petri dishes (4-5 mm thickness). Bacterial suspensions of *Staphylococcus aureus* and *Escherichia coli* (100 μ L) were lawned on separate MHA plates. Four wells were punched per plate and filled with: (1) probiotic hydrogel, (2) probiotic cream, (3) *L. paracasei* supernatant, and (4) negative control. Plates were incubated at 37°C for 24 and 48 hours. Zones of inhibition were measured using a vernier caliper and interpreted as: strong (>20 mm), intermediate (10-20 mm), or weak (<10 mm).

2.4. *In Vivo* Wound Healing Assay of *Postbiotic*-formulated Hydrogel

Thirty healthy male ICR mice (*Mus musculus*), eight weeks of age and weighing 25-30 grams at the start of the study, were obtained from the Laboratory Animal Care Facility of Mariano Marcos State University (MMSU), Ilocos Norte, Philippines. Animals were housed individually under controlled environmental conditions (22-25 °C; 55-75% relative humidity; 12 hours light/dark cycle) and were provided standard laboratory chow and water *ad libitum*. A five-day acclimatization period was observed

prior to the initiation of experimental procedures. After acclimatization, twenty-four mice were subjected to full-thickness excisional wounds created in the dorso-lumbar region along the midline (vertebral levels L1-L6) using a sterile 1.0 cm biopsy punch. All procedures were performed under topical lidocaine anesthesia. The remaining six mice were maintained as unwounded basal controls.

Treatments commenced 24 hours after wound induction and were administered once daily using sterile applicators, with formulations applied to cover the wound bed and extend approximately 5 mm beyond the wound margins. Animals were assigned to the following groups: (1) experimental group treated with *L. paracasei* 7060-formulated nipa hydrogel; (2) negative control group receiving hydrogel base only; (3) positive control group treated with DuoDERM hydrogel supplemented with povidone-iodine; (4) untreated negative control group; and (5) basal group (unwounded).

Wound size and gross morphological features were monitored at designated time points. Tissue samples were collected on days 3 and 21 ($n = 3$ per group per time point) for gene expression analysis. All animal procedures were reviewed and approved by the Institutional Animal Care and Use Committee (IACUC) of Mariano Marcos State University and were conducted in compliance with institutional and national guidelines for the ethical use of laboratory animals.

2.5. Gene Expression Analysis

Gene-specific primers for selected wound healing markers were designed using the Integrated DNA Technologies PrimerQuest tool, based on genomic sequences from the NCBI database. All primers were validated for specificity and amplicon size prior to use in qPCR. Total RNA was extracted from wound tissues using TRIzol Reagent (Invitrogen) and quantified with a ScanDrop2 spectrophotometer (Analytik Jena). Two micrograms of RNA per sample were treated with RQ1 RNase-Free DNase I (Vivantis) to eliminate genomic DNA. qRT-PCR was performed using the SuperScript™ III Platinum™ One-Step qRT-PCR Kit (Invitrogen) on a QuantStudio™ 5 Real-Time PCR System (Applied Biosystems). Thermal cycling was conducted at 95°C for 30 seconds, followed by 39 cycles of 95°C for 5 seconds, 60°C for 30 seconds, and 72°C for 30 seconds. Melt curve

analysis confirmed amplification specificity. Relative gene expression was normalized to GAPDH using the ΔCt method. Samples were collected on days 3 and 21 post-wounding, corresponding to the inflammatory and remodeling phases, respectively (Table 1).

2.6. Whole Genome Sequencing

Genomic DNA was extracted from *L. paracasei* 7060 using the Zymo Quick-DNA™ Fungal/Bacterial Miniprep Kit (Cat. No. D6005), with mechanical disruption performed on the MP Biomedicals FastPrep 96™ system to enhance cell lysis efficiency. DNA quality and yield were assessed using the LightBench® platform, which confirmed the presence of high-molecular-weight DNA at concentrations suitable for long-read sequencing.

High-molecular-weight DNA was subsequently sheared and size-selected following the PacBio SMRTbell library preparation guidelines. End-repair and adapter ligation steps were carried out to generate SMRTbell libraries. Sequencing was performed on the PacBio Sequel IIe system, producing long-read data with read lengths exceeding 10 kb.

Raw subreads were assembled using the Hierarchical Genome Assembly Process (HGAP), resulting in a highly contiguous draft genome assembly. The assembled contigs were then submitted to the Rapid Annotation using Subsystem Technology (RAST) server for automated genome annotation, enabling the assignment of putative functions and subsystem classifications to predicted coding sequences.

2.7. Promoter Prediction and Transcription Factor Binding Site Analysis

Putative promoter elements upstream of selected genes were identified using *in silico* analysis. Upstream regions (approximately 250 bp) were extracted from the assembled whole genome sequence of *L. paracasei* 7060. Promoter motifs, including -10 and -35 elements, were predicted using the BPROM tool (Softberry, Inc.), which provided positional information and predicted Transcription Start Sites (TSS). To further identify potential transcription factor binding sites, sequences were analyzed using the Virtual Footprint tool from the PRODORIC database, which screens for conserved regulatory motifs across known bacterial transcription factor binding profiles.

Table 1. Primer sets used for quantitative gene expression analysis.

Gene and ID	Species	Primer Sequence(5' - 3')	Product Size(bp)
<i>TNF-α</i> 21926	<i>Mus musculus</i> L.	F- TTGTCTACTCCAGGTTCTCT	107
		R- GAGGTTGACTTTCTCCTGGTATG3	
<i>TGF-β</i> 21803	<i>Mus musculus</i> L.	F- CCTGCAAGACCATCGACAT	90
		R- GACTGGCGAGCCTTAGTTT	
<i>TLR2</i> 24088	<i>Mus musculus</i> L.	F- CACTATCCGGAGGTTGCATATC	113
		R- GGAAGACCTTGCTGTTCTCTAC	

2.8. Histological and Morphometric Assessments

Wound tissues were evaluated histologically using a semi-quantitative grading system adapted from Gupta and Kumar [24]. The assessment focused on key parameters of wound healing, including granulation tissue formation, degree of inflammatory infiltrate, collagen fiber orientation, collagen pattern, and the deposition of early and mature collagen (Table 2, reproduced from Gupta A, Kumar P. *Assessment of the histological state of the healing wound*, under the terms of the Creative Commons Attribution 4.0 International License [24]). Histological scoring was performed in a blinded manner to minimize observer bias.

Table 2. Histological grading system for assessment of wound healing.

Number	Histological Parameters
1	Amount of granulation tissue (profound-1, moderate-2, scanty-3, absent-4)
2	Inflammatory infiltrate (plenty-1, moderate-2, a few-3)
3	Collagen fiber orientation (vertical-1, mixed-2, horizontal-3)
4	Pattern of collagen (reticular-1, mixed-2, fascicle-3)
5	Amount of early collagen (profound-1, moderate-2, minimal-3, absent-4)
6	Amount of mature collagen (profound-1, moderate-2, minimal-3)

Macroscopic wound appearance was graded based on established qualitative criteria. Scores ranged from A (very red, swollen, moist) to F (hair regrowth and complete wound closure).

Digital images of wounds were captured every 24 hours post-treatment. Wound area was measured using ImageJ® software (v1.54p), calibrated with a ruler. Wound contraction rate (%) was computed as (Eq. 1):

Healing Rate

$$= \frac{\text{Initial Wound Diameter (mm)} - \text{Final Wound Diameter (mm)}}{\text{Initial Wound Diameter (mm)}} \times 100 \quad (1)$$

Red, green, and blue (RGB) values of wound areas were extracted from standardized images using ImageJ®, with normalization against a reference image to control for lighting variability. Chromatic red (r) was calculated as (Eq. 2), and luminance (L) was calculated as (Eq. 3):

$$r = \frac{(100 \times R)}{R+G+B} \quad (2)$$

$$L = \frac{R+G+B}{3} \quad (3)$$

2.9. Data Analysis

Statistical differences among treatment groups were assessed using one-way ANOVA. When significant F-values were observed, Tukey's Honest Significant Difference (HSD) test was applied for post hoc comparisons. Analyses were performed using SAS software (v9.4; SAS Institute Inc.).

2.10. Ethical Considerations

All procedures involving animals were approved by the Institutional Animal Care and Use Committee (IACUC) of Mariano Marcos State University and conducted in accordance with the ethical guidelines of the National Committee for Research Ethics in Science and Technology (NENT, 2018). Efforts were made to minimize animal suffering and ensure humane care throughout the study.

3. RESULTS

3.1. Morphoanatomical Changes in the Wound under Postbiotic Hydrogel

On day 0, all wounds were of similar size (~10 mm) and exhibited bright red color, characteristic of acute injury. By day 3, the *L. paracasei* 7060 hydrogel-treated wounds (iv) and DuoDerm-treated wounds (iii) showed early signs of fibrin clotting and color shift to yellow-brown, indicating reduced inflammation. By days 6–9, these same wounds showed significant contraction, scab darkening, and drying, which are morphological hallmarks of wound healing and epithelialization. Full closure was observed in group (iv) by day 12, followed by fur regrowth and minimal scarring between days 15–21. In contrast, groups treated with base hydrogel (ii) and DuoDerm + povidone-iodine (iii) healed more slowly, with persistent scabbing and incomplete closure by day 15. Untreated wounds (i) showed the poorest recovery: open wounds with little fur regrowth even on day 21 (Fig. 1a). Wound contraction was fastest in the *L. paracasei* 7060 hydrogel and DuoDerm groups, reaching 60–70% reduction by day 9 and complete closure by day 15. Other groups retained more than 50% of wound area by day 9 and showed incomplete healing at day 21 (Fig. 1b).

Color changes were most evident in the *L. paracasei* hydrogel group (iv). Over the course of healing, these wounds gradually transitioned from a dark reddish-brown appearance to lighter pink hues. In comparison, wounds in the untreated group (i) largely retained darker coloration throughout the observation period, consistent with sustained inflammation or slower remodeling (Fig. 2a). The Chromatic Red Index (CRI) decreased in all treatment groups; however, the most pronounced reduction was observed in group ii between days 6 and 15 (Fig. 2b, top). Luminance values (Fig. 2b, bottom) showed a marked increase in this group after day 15, reflecting increased wound brightness and suggesting ongoing re-epithelialization. In contrast, the DuoDerm (iii) and base hydrogel (ii) treatments exhibited only moderate reductions in CRI and comparatively smaller changes in luminance. Tukey's post hoc analysis confirmed significant differences in both CRI and luminance among treatment groups at several time points, with the strongest effects detected after day 6 ($p \leq 0.01$).

3.2. Histological Evaluation of Wound Healing

Histological examination of wound sections harvested on day 21 revealed clear differences in tissue organization and inflammatory status among the treatment groups

(Fig. 3a). Healing scores were determined using the semi-quantitative histological grading criteria outlined in Table 2. Wounds treated with the base hydrogel showed pronounced inflammatory cell infiltration and poorly organized granulation tissue. Collagen fibers in this group were predominantly immature and vertically oriented, reflecting limited matrix remodeling. Accordingly, this group received a healing score of 4, consistent with suboptimal tissue regeneration.

In comparison, wounds treated with Duoderm combined with povidone-iodine displayed moderate improvements in tissue architecture. These sections exhibited partial re-epithelization, reduced inflammatory infiltrates, and early signs of extracellular matrix organization, corresponding to a healing score of 8. Interestingly, wounds from the *L. paracasei*-treated group and the untreated group demonstrated largely comparable histological features.

Both groups showed a well-reestablished epidermal layer, evidence of hair follicle regeneration, and minimal residual inflammation, indicative of advanced wound repair. As a result, both groups received high healing scores of 12, reflecting favorable tissue recovery by day 21. Detailed semi-quantitative lesion scores are provided in **Supplementary A**.

Despite the observed histological improvements associated with probiotic treatment, agar diffusion assays indicated that none of the hydrogel formulations, including the *L. paracasei* 7060 hydrogel, exhibited detectable antimicrobial activity against *Staphylococcus aureus*, *Escherichia coli*, or *Candida albicans*. In contrast, penicillin-treated positive controls produced distinct zones of inhibition against *S. aureus* and *C. albicans* (>20mm), whereas *E. coli* remained resistant to all treatments, including penicillin (Fig. 3b).




Fig. (1). Epithelial tissue repair is accelerated when the wound is treated with a probiotic gel containing *L. paracasei* 7060. **a)** Representative photographs of epithelial tissue repair in different treatments. Each column represents a treatment which includes i) untreated, ii) hydrogel base, iii) Duoderm hydrogel with povidone-iodine, and iv) postbiotic hydrogel. **b)** Comparative wound size and wound contraction of the different treatments in the 21-day observation period. Statistical differences were assessed using ANOVA and Tukey's multiple comparison test at $\alpha = 0.05$. Error bars indicate the standard error calculated from biological triplicate. The condition was found to be a significant source of variation at $p \leq 0.01$.




Fig. (2). Epithelial tissue repair is accelerated when the wound is treated with a probiotic gel containing *L. paracasei* 7060. **a)** Visual representation of color change in wound from day 3 to day 21 (rows) across four treatment groups (i-iv, columns). Each square represents the mean color appearance of replicates at a given time point. Each column represents a treatment which includes i) untreated, ii) hydrogel base, iii) DuoDerm hydrogel, and iv) postbiotic hydrogel. **b)** Quantitative assessment of color parameters: Top panel, Chromatic Red Index values showing a general decline in redness across all treatments with increasing storage time. Bottom panel, Luminance values indicating a progressive lightening of samples over time, particularly after day 15. Statistical differences were assessed using ANOVA and Tukey's multiple comparison test at $\alpha = 0.05$. Error bars indicate the standard error calculated from biological triplicate. The condition was found to be a significant source of variation at $p \leq 0.01$.

3.3. Differential Expression of Tissue-repair Associated Genes

The relative expression levels of three key genes involved in inflammation and tissue repair, *TLR2*, *TNF- α* , and *TGF- β* , were assessed in ICR mice on days 3 and 21 post-wounding across four treatment groups: Postbiotic Hydrogel, Hydrogel Base, DuoDerm Hydrogel, and Untreated Control (Fig. 4a).

At day 3, *TNF- α* expression was significantly elevated across all experimental groups relative to baseline levels. The hydrogel base group exhibited the highest *TNF- α* expression, surpassing even that observed in the untreated control. In contrast, the postbiotic hydrogel-treated group showed substantially lower *TNF- α* and *TGF- β* expression

compared with the hydrogel base group, with *TNF- α* levels falling below those of the untreated control. *TLR2* expression was upregulated in all treatment groups at day 3, with the most pronounced increase detected in the postbiotic hydrogel group.

By day 21, expression levels of *TNF- α* and *TLR2* had decreased markedly in most groups, consistent with progression beyond the acute inflammatory phase. The most pronounced reduction in *TLR2* expression was observed in the postbiotic hydrogel group, whereas wounds treated with DuoDERM® hydrogel maintained comparatively higher *TLR2* levels. In contrast, *TGF- β* expression remained at moderate levels in the postbiotic hydrogel group, while more substantial downregulation was observed in the other treatment groups.




Fig. (3). Histological characteristics of epithelial tissues in the dorso-lumbar (midline over L1-L6) 21 days post-treatment and antimicrobial effects of postbiotic hydrogel treatment. **a)** Representative photomicrographs of H&E-stained histological sections showing epithelial tissue repair across treatment groups: i) postbiotic hydrogel, ii) hydrogel base, iii) DuoDerm hydrogel with povidone-iodine, and iv) untreated control. **b)** antimicrobial activity of treatments against *Staphylococcus aureus*, *Escherichia coli*, and *Candida albicans*. Top panel: Representative images of disk diffusion assay; bottom panel: quantitative assessment of the inhibition zones. Statistical differences were assessed using ANOVA and Tukey's multiple comparison test at $\alpha = 0.05$. Error bars indicate the standard error calculated from biological triplicate. The condition was found to be a significant source of variation at $p \leq 0.01$.

At three days post-wounding, *L. paracasei* 7060 demonstrated a modulatory effect on the inflammatory response, facilitating the transition toward the early proliferative phase of wound healing. As illustrated in Fig. (4b), *L. paracasei* 7060 was shown to interact with *TLR2*, leading to pronounced *TLR2* activation, a recognized indicator of innate immune engagement. This interaction was associated with downstream activation of the *NF- κ B* signaling pathway, which subsequently supported the transcription of immune response-related genes. Activation of this pathway did not translate into excessive inflammatory signaling. Instead, *TNF- α* expression appeared to be tightly regulated, suggesting a controlled pro-inflammatory response that is conducive to effective tissue repair rather than sustained tissue damage. At the same time, an increase in *TGF- β* expression was observed, consistent with the initiation of the transition from the inflammatory phase toward the early proliferative stage of wound healing. These molecular signatures are indicative of a controlled immune modulation by *L. paracasei* 7060, supporting its role in enhancing wound healing dynamics (Fig. 4b).

3.4. Whole Genome Sequence (WGS) Analysis of *L. paracasei* 7060

Whole genome analysis of *L. paracasei* 7060 revealed a diverse set of genes linked to the production of bioactive metabolites and cellular components involved in antimicrobial activity, immune modulation, and tissue repair (Fig. 5b). In particular, the genome contained biosynthetic gene clusters encoding bacteriocins, lactic acid, Short-Chain Fatty Acids (SCFAs), NADH, and Lipoteichoic Acid (LTA) (Fig. 5a and 5b). The identification of bacteriocin-related genes supports the strain's capacity to produce antimicrobial peptides that may contribute to the inhibition of pathogenic bacteria. Genes associated with lactate and acetate biosynthesis were also detected, indicating a potential role in promoting angiogenesis and reinforcing epithelial barrier function, respectively. In addition, the presence of NADH-producing genes suggests a contribution to cellular redox balance and mitigation of oxidative stress within the wound microenvironment. Genes involved in lipoteichoic acid biosynthesis were likewise identified and are known to participate in host immune modulation through *TLR2*-mediated signaling.





Fig. (4). Gene expression and involvement in tissue repair. **a)** Relative gene expression levels (log₂ fold change) of pro- and anti-inflammatory markers: *Tumor Necrosis Factor alpha* (*TNF-α*), *Transforming Growth Factor beta* (*TGF-β*), and *Toll-Like Receptor 2* (*TLR2*) in treated wounds. The top panel shows upregulation; the bottom panel shows downregulation of gene expression across treatment groups. **b)** Schematic representation of *L. paracasei* 7060-mediated modulation of the inflammatory to early proliferative transition phase at 3 days post-wounding. Statistical differences were assessed using ANOVA and Tukey's multiple comparison test at $\alpha = 0.05$. Error bars indicate the standard error calculated from biological triplicate. The condition was found to be a significant source of variation at $p \leq 0.01$.



b)

Category	Map ID	Genes and Transcription Factors	Function	References
Immunity	JJMPBGNO_04407	Carnobacteriocin-B2 immunity Protein	acts as an antimicrobial peptide	Quadri L et al, 1997
Tissue Repair and Angiogenesis	JJMPBGNO_04918	L-lactate dehydrogenase	stimulates collagen and vascular endothelial growth factor	Ghani Q.P. et al, 2004
	JJMPBGNO_01866			
	JJMPBGNO_01959	Lactate 2-monoxygenase	Tissue repair and formation of new blood vessels	Hunt TK et al, 2007 Manosalva et al, 2022
	JJMPBGNO_02537			
	JJMPBGNO_00256			
	JJMPBGNO_02422			
	JJMPBGNO_02421	Lactate utilization protein A	Stimulates the release of angiogenic factors like VEGF and fibroblasts for collagen	Lee TY, 2021 Huiting W., 2023
	JJMPBGNO_02421	Lactate utilization protein B	promotes the proliferation of macrophages	Ghani Q.P. et al, 2004
	JJMPBGNO_03968	Lactate racemase	promoting efferocytosis-induces macrophage proliferation	Nature Metabolism 2023 Hunt TK et al, 2007
	JJMPBGNO_03968	Lactate racemase	initiates connective tissue synthesis	Fang Y et al, 2004
JJMPBGNO_03969	Lactate racemization regulatory protein	amplify the inflammatory response; retention of CD4+ and CD8+ T cells	Zitong W et al 2024 Wang Z et al, 2023	
JJMPBGNO_01208	NADH oxidase	Promotes the expression of genes linked to tissue repair	Huiting W., 2023	
JJMPBGNO_01248	NADH oxidase	Influences immune regulation, inflammation control, and proliferative remodeling	Chan E.C. et al, 2014	
JJMPBGNO_02733	Lipoteichoic acid synthase 1	Influences the production and remodeling of the extracellular matrix	Chan E.C. et al, 2014	
JJMPBGNO_03917	Lipoteichoic acid synthase 1	promoting cell proliferation, angiogenesis, and collagen synthesis	Levigne D, 2016	
JJMPBGNO_02733	Lipoteichoic acid synthase 1	Induce epithelial-mesenchymal transition and wound healing in HaCaT cells	Oku Y et al, 2009	
Metabolic Regulation / Remodeling	JJMPBGNO_01250	Acetate kinase	metabolic homeostasis	Yoshida Y et al, 2019
	JJMPBGNO_01918			
	JJMPBGNO_04802			
Stress Response & Survival	JJMPBGNO_04533	LexA	helps maintain probiotic function during host stress conditions	Dalbanjan et al, 2024
	JJMPBGNO_03054	Peroxide-responsive PerR	increases bacterial resilience in inflammatory wounds with high H ₂ O ₂ levels	Vliet A & Janssen-Heininger YM, 2014
	JJMPBGNO_00466	Heat-inducible transcription HrcA	Helps bacteria withstand host-induced stress (heat, oxidative)	Hakim et al, 2020
	JJMPBGNO_00657	Redox-sensing transcriptional Rex	Supports bacterial survival under oxidative stress, maintaining colonization	Zhu H et al, 2018

Fig. (5). (a) Circular genome map of *L. paracasei* 7060. The outermost rings represent predicted coding sequences (CDSs) on the forward (outermost, purple) and reverse (second outermost, purple) strands. The next two inner rings (green) display the locations of rRNA and tRNA genes. The blue and orange rings show the GC content and GC skew. (b) Functional classification of genes and transcription factors in *L. paracasei* 7060 associated with the biosynthesis of metabolites potentially involved in tissue repair.

Beyond metabolic functions, the genome also harbored genes associated with stress adaptation and host interaction. Among these, *trxA*, encoding thioredoxin, is implicated in redox regulation and protection against Reactive Oxygen Species (ROS). The detection of *msoR* and *gshA*, which are involved in metal sensing and glutathione biosynthesis, respectively, further points to roles in detoxification processes and epithelial protection. Genes encoding the *spaCBA* pilus cluster were present, suggesting a potential mechanism for adhesion to host epithelial surfaces. Multiple copies of *ef-Tu* were also observed; this protein is recognized for its moonlighting functions, including interactions with host pattern recognition receptors and modulation of immune responses. The genome map (Fig. 5a) illustrates the circular architecture of the *L. paracasei* 7060 chromosome, highlighting coding sequences (CDSs), rRNA and tRNA loci, as well as regions of GC content and GC skew.

4. DISCUSSION

The integration of phenotypic, histological, gene expression, and genomic data in this study provides converging evidence supporting the pro-regenerative potential of a postbiotic hydrogel formulated with *Lactobacillus paracasei* 7060. The findings indicate that treatment with this hydrogel markedly accelerated tissue repair, as demonstrated by complete wound closure by day 7 and full re-epithelialization by day 21. These outcomes suggest that microbial-derived metabolites present in the postbiotic formulation actively modulate host immune responses and repair mechanisms rather than merely providing passive wound coverage. Similar regenerative effects have been reported for probiotics and their postbiotic derivatives, which have been shown to promote epithelial regeneration, enhance collagen deposition, and regulate inflammatory responses during wound healing [25-27].

In the present study, wounds treated with the *L. paracasei* 7060 postbiotic hydrogel exhibited clear improvements in surface characteristics, including color, luminance, homogeneity, smoothness, and consistency. These parameters are widely recognized as indicators of accelerated and high-quality tissue repair [28]. By day 21, complete re-epithelialization was accompanied by histological evidence of increased collagen deposition, likely involving collagen types I and III, as well as enhanced angiogenesis [29]. Both collagen remodeling and neovascularization are central to wound maturation and are essential for restoring tissue integrity, tensile strength, and long-term functionality [30, 31]. Phenotypic differences observed at earlier stages were not reflected histologically by day 2 [32], suggesting that the most dynamic phase of tissue regeneration may have occurred earlier in the healing timeline, potentially around day 12 [33,34]. This interpretation is consistent with previous reports indicating that, by day 21, wound resolution is generally underway even in untreated wounds, although the quality of healing, including scar formation and tissue organization, remains strongly influenced by early

inflammatory and reparative events [35, 36].

At the molecular level, the postbiotic hydrogel induced coordinated changes in the expression of genes involved in inflammation and tissue remodeling. Early downregulation of the pro-inflammatory cytokine TNF- α and the pro-fibrotic mediator TGF- β suggests attenuation of excessive inflammatory signaling during the initial repair phase. TNF- α is a key driver of acute inflammation, and its suppression is associated with reduced leukocyte infiltration, diminished secondary tissue damage, and improved healing outcomes [37, 38]. Likewise, although TGF- β plays a critical role in extracellular matrix deposition and wound remodeling, sustained or excessive expression of this cytokine can promote pathological fibrosis and impair normal tissue regeneration [39]. The reduced expression of both TNF- α and TGF- β early in the repair process, therefore, points toward a healing trajectory that favors regeneration over fibrosis, ultimately contributing to improved tissue architecture and function [40].

In contrast, Toll-Like Receptor 2 (TLR2) expression was significantly upregulated during the early stages of healing, followed by a gradual decline at later time points. Activation of TLR2 by microbial ligands, including lipoteichoic acids derived from Gram-positive bacteria, is known to stimulate innate immune responses and support tissue repair when appropriately regulated [41-43]. The observed pattern of early induction followed by controlled downregulation may have created a transient window of immune activation sufficient to promote pathogen clearance, macrophage recruitment, and early repair signaling, while preventing prolonged inflammation. Such temporal regulation of TLR signaling is increasingly recognized as a hallmark of optimal wound healing responses [44]. The coordinated dynamics observed between TLR2, TNF- α , and TGF- β suggest that the postbiotic hydrogel fosters an immunological environment conducive to rapid inflammation resolution and effective tissue remodeling, potentially minimizing long-term scarring. This profile aligns well with current wound healing paradigms emphasizing early inflammatory control, timely macrophage polarization, and structured extracellular matrix reorganization [45].

Genomic characterization of *L. paracasei* 7060 further supports the biological plausibility of these observed effects. The genome harbors genes encoding enzymes involved in the production of lactic acid and Short-Chain Fatty Acids (SCFAs), including acetate, metabolites with well-established roles in inflammation modulation and tissue regeneration. Lactate has been shown to promote angiogenesis by stimulating endothelial cell proliferation and Vascular Endothelial Growth Factor (VEGF) expression [46], while acetate contributes to epithelial barrier reinforcement and immune homeostasis through signaling pathways involving G-protein-coupled receptors such as GPR43 [47]. In addition, the presence of genes linked to NADH biosynthesis suggests a capacity for antioxidant support, which is particularly relevant in the highly oxidative environment of acute and chronic wounds. NADH functions as a critical cofactor for glutathione

reductase and other redox enzymes, helping to counteract Reactive Oxygen Species (ROS)-mediated damage and maintain cellular viability during tissue repair [48, 49].

Genes associated with lipoteichoic acid (LTA) biosynthesis were also identified, providing a potential molecular basis for the observed modulation of TLR2 signaling [50]. Controlled interactions between LTA and TLR2 are known to activate beneficial immune pathways without triggering excessive inflammation, thereby supporting macrophage-mediated tissue repair processes [51]. Furthermore, the presence of antioxidant-related genes such as *trxA*, *msoR*, and *gshA*, encoding components of the glutathione system, underscores the strain's potential to maintain redox balance and cellular homeostasis under stress conditions, which are factors that are critical not only for microbial resilience but also for epithelial healing [52, 53].

The identification of the *spaCBA* gene cluster, which encodes pilus-associated proteins involved in adhesion, suggests that close microbial-host interactions may facilitate localized delivery of bioactive metabolites and enhance targeted immune modulation at epithelial surfaces [54]. Additionally, the presence of multiple copies of *ef-Tu* introduces another layer of immunomodulatory potential, as this moonlighting protein has been reported to interact with host pattern recognition receptors and influence cytokine expression profiles [55, 56]. Taken together, these genomic features provide a mechanistic framework explaining how postbiotics derived from *L. paracasei* 7060 may orchestrate a multifaceted regenerative response.

The findings of this study demonstrate that *L. paracasei* 7060-derived postbiotics promote wound healing through coordinated suppression of excessive early inflammation, enhancement of antioxidant defenses, stimulation of angiogenesis, and support of epithelial regeneration. The integration of phenotypic observations with molecular and genomic data strengthens the case for this strain as a promising candidate for future wound care applications, including hydrogel-based delivery systems, tissue-engineering scaffolds, or topical formulations. Future investigations should aim to validate these effects in large animal models and human skin equivalents, as well as to delineate the individual contributions of specific microbial metabolites to tissue repair. Longitudinal analyses of wound-associated microbiome dynamics and immune cell phenotypes would further advance understanding of the complex host-microbe interactions that shape wound healing outcomes.

CONCLUSION

This study provides evidence for the wound healing potential of a postbiotic hydrogel derived from *L. paracasei* 7060. Treatment with the formulation accelerated wound closure, enhanced collagen deposition, and promoted complete re-epithelialization, reflecting an optimized balance between inflammation and regeneration. The coordinated molecular and phenotypic outcomes highlight *L. paracasei* 7060 as a promising source of bioactive postbiotics for wound healing applications. Further studies using advanced preclinical models will be important to

confirm these effects and to clarify the specific roles of individual microbial metabolites in driving tissue regeneration.

AUTHORS' CONTRIBUTIONS

The authors confirm their contribution to the paper as follows: A.G.B.P., J.Z.P.F., M.J.T.T.A., A.D.: Methodology; A.G.B.P.: Data collection; A.G.B.P., J.Z.P.F., J.P.A.A.: Analysis and interpretation of results; A.G.B.P., J.Z.P.F., P.J.I.G.: Draft manuscript. All authors reviewed the results and approved the final version of the manuscript.

LIST OF ABBREVIATIONS

ANOVA	= Analysis of Variance
bp	= Base Pair
cDNA	= Complementary DNA
CMC	= Carboxymethylcellulose
CRD	= Completely Randomized Design
CRI	= Chromatic Red Index
DNA	= Deoxyribonucleic Acid
ECM	= Extracellular Matrix
ef-Tu	= Elongation Factor Tu
GAPDH	= Glyceraldehyde-3-Phosphate Dehydrogenase
GC	= Guanine-Cytosine
gsha	= Glutathione Synthetase Gene
H&E	= Hematoxylin and Eosin
HSD	= Honest Significant Difference
IACUC	= Institutional Animal Care and Use Committee
IDT	= Integrated DNA Technologies
IL	= Interleukin
ISAPP	= International Scientific Association for Probiotics and Prebiotics
LAB	= Lactic Acid Bacteria
LTA	= Lipoteichoic Acid
MHA	= Mueller-Hinton Agar
MRS	= de Man, Rogosa, and Sharpe (culture medium)
msor	= Metal-Sensing Oxidoreductase Gene
NCBI	= National Center for Biotechnology Information
NF- κ b	= Nuclear Factor Kappa B
NENT	= National Committee for Research Ethics in Science and Technology
PBS	= Phosphate-Buffered Saline
qPCR	= Quantitative Polymerase Chain Reaction

RAST	= Rapid Annotation using Subsystem Technology
RGB	= Red-Green-Blue (color analysis)
RNA	= Ribonucleic Acid
ROS	= Reactive Oxygen Species
RPM	= Revolutions Per Minute
SCFA	= Short-Chain Fatty Acid
SMRTBELL	= Single Molecule Real-Time Bell (PacBio library format)
TGF- β	= Transforming Growth Factor-Beta
TLR2	= Toll-Like Receptor 2
TNF- α	= Tumor Necrosis Factor-Alpha
TSS	= Transcription Start Site
trxa	= Thioredoxin
VEGF	= Vascular Endothelial Growth Factor
WGS	= Whole Genome Sequencing

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

This study was approved by the Institutional Animal Care and Use Committee (IACUC) of Mariano Marcos State University under protocol number 2024-001. In addition, animal research clearance was secured from the Bureau of Animal Industry of the Philippines under reference number AR-2024-0505.

HUMAN AND ANIMAL RIGHTS

This study adheres to internationally accepted standards for animal research, following the 3Rs principle. The ARRIVE guidelines were employed for reporting experiments involving live animals, promoting ethical research practices.

All procedures involving animals were conducted in accordance with the ethical guidelines of the National Committee for Research Ethics in Science and Technology (NENT, 2018).

CONSENT FOR PUBLICATION

Not applicable.

AVAILABILITY OF DATA AND MATERIALS

All data generated or analyzed during this study are included in this published article.

FUNDING

This study was supported by the Commission on Higher Education (CHED), Philippines, with project number LAKAS 2022-005.

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

The authors are thankful to the CHED Philippines for supporting the project. The authors are also thankful to Dr. Shirley C. Agrupis, Dr. Dionisio S. Bucaco, and Dr. Rhian Jaymar D. Ramil for their administrative support.

SUPPLEMENTARY MATERIAL

Supplementary material is available on the publisher's website along with the published article.

REFERENCES

- [1] Marchiando AM, Graham WV, Turner JR. Epithelial barriers in homeostasis and disease. *Annu Rev Pathol* 2010; 5(1): 119-44. <http://dx.doi.org/10.1146/annurev.pathol.4.110807.092135> PMID: 20078218
- [2] Gurtner GC, Werner S, Barrandon Y, Longaker MT. Wound repair and regeneration. *Nature* 2008; 453(7193): 314-21. <http://dx.doi.org/10.1038/nature07039> PMID: 18480812
- [3] Ruffin M, Brochiero E. Repair process impairment by *Pseudomonas aeruginosa* in epithelial tissues: Major features and potential therapeutic avenues. *Front Cell Infect Microbiol* 2019; 9(182): 182. <http://dx.doi.org/10.3389/fcimb.2019.00182> PMID: 31214514
- [4] Lauterio M, Deck DH. Current challenges in the management of skin and soft tissue infections and community-acquired pneumonia. *J Fam Pract* 2022; 71(5 Suppl): S2-9. <http://dx.doi.org/10.12788/jfp.0423> PMID: 35776861
- [5] Wang J, Chao J. Epithelial cell dysfunction in pulmonary fibrosis: Mechanisms, interactions, and emerging therapeutic targets. *Pharmaceuticals* 2025; 18(6): 812. <http://dx.doi.org/10.3390/ph18060812> PMID: 40573209
- [6] Klingberg F, Hinz B, White ES. The myofibroblast matrix: Implications for tissue repair and fibrosis. *J Pathol* 2013; 229(2): 298-309. <http://dx.doi.org/10.1002/path.4104> PMID: 22996908
- [7] Oelschlaeger TA. Mechanisms of probiotic actions - A review. *Int J Med Microbiol* 2010; 300(1): 57-62. <http://dx.doi.org/10.1016/j.ijmm.2009.08.005> PMID: 19783474
- [8] Yin Z, Wang Y, Feng X, et al. *Lactobacillus rhamnosus*GG and *Bifidobacterium animalis* subsp. *lactis*BB-12 promote infected wound healing via regulation of the wound microenvironment. *Microb Biotechnol* 2024; 17(10): e70031. <http://dx.doi.org/10.1111/1751-7915.70031> PMID: 39422648
- [9] Saravanan P, R P, Balachander N, K KRS, S S, S R. Anti-inflammatory and wound healing properties of lactic acid bacteria and its peptides. *Folia Microbiol* 2023; 68(3): 337-53. <http://dx.doi.org/10.1007/s12223-022-01030-y> PMID: 36780113
- [10] Knackstedt R, Knackstedt T, Gatherwright J. The role of topical probiotics in skin conditions: A systematic review of animal and human studies and implications for future therapies. *Exp Dermatol* 2020; 29(1): 15-21. <http://dx.doi.org/10.1111/exd.14032> PMID: 31494971
- [11] Kong Y, Olejar KJ, On SLW, Chelikani V. The potential of *Lactobacillus* spp. for modulating oxidative stress in the gastrointestinal tract. *Antioxidants* 2020; 9(7): 610. <http://dx.doi.org/10.3390/antiox9070610> PMID: 32664392
- [12] Aguilar-Toalá JE, Garcia-Varela R, Garcia HS, et al. Postbiotics: An evolving term within the functional foods field. *Trends Food Sci Technol* 2018; 75: 105-14. <http://dx.doi.org/10.1016/j.tifs.2018.03.009>
- [13] Salminen S, Collado MC, Endo A, et al. The International Scientific Association of Probiotics and Prebiotics (ISAPP) consensus statement on the definition and scope of postbiotics. *Nat Rev Gastroenterol Hepatol* 2021; 18(9): 649-67. <http://dx.doi.org/10.1038/s41575-021-00440-6> PMID: 33948025
- [14] Tsilingiri K, Rescigno M. Postbiotics: What else? *Benef Microbes*

- 2013; 4(1): 101-7.
<http://dx.doi.org/10.3920/BM2012.0046> PMID: 23271068
- [15] Ulluwishewa D, Anderson RC, McNabb WC, Moughan PJ, Wells JM, Roy NC. Regulation of tight junction permeability by intestinal bacteria and dietary components. *J Nutr* 2011; 141(5): 769-76.
<http://dx.doi.org/10.3945/jn.110.135657> PMID: 21430248
- [16] Schultz GS, Chin GA, Moldawer L, Diegelmann RF. Principles of wound healing. In: Fitridge R, Thompson M, Eds. *Mechanisms of vascular disease: A reference book for vascular specialists*. Adelaide, Australia: University of Adelaide Press 2011; pp. 481-516.
<http://dx.doi.org/10.1017/UPO9781922064004.024>
- [17] López P, González-Rodríguez I, Gueimonde M, Margolles A, Suárez A. Immune response to *Bifidobacterium bifidum* strains support Treg/Th17 plasticity. *PLoS One* 2011; 6(9): e24776.
<http://dx.doi.org/10.1371/journal.pone.0024776> PMID: 21966367
- [18] Arqués JL, Rodríguez E, Langa S, Landete JM, Medina M. Antimicrobial activity of lactic acid bacteria in dairy products and gut: Effect on pathogens. *BioMed Res Int* 2015; 2015: 1-9.
<http://dx.doi.org/10.1155/2015/584183> PMID: 25861634
- [19] Umekar M, Chaudhary AA, Manghani M, et al. Probiotics in nanotechnology-driven wound healing: From mechanistic insight to clinical promise. *Pharmaceutics* 2025; 17(7): 805.
<http://dx.doi.org/10.3390/pharmaceutics17070805> PMID: 40733015
- [20] Xie A, Chen A, Chen Y, et al. *Lactobacillus* for the treatment and prevention of atopic dermatitis: Clinical and experimental evidence. *Front Cell Infect Microbiol* 2023; 13: 1137275.
<http://dx.doi.org/10.3389/fcimb.2023.1137275> PMID: 36875529
- [21] Alves AC, Martins SMSB Jr, Belo JVT, et al. Global trends and scientific impact of topical probiotics in dermatological treatment and skincare. *Microorganisms* 2024; 12(10): 2010.
<http://dx.doi.org/10.3390/microorganisms12102010> PMID: 39458319
- [22] Machado P, Ribeiro FN, Giublin FCW, et al. Next-generation wound care: A scoping review on probiotic, prebiotic, synbiotic, and postbiotic cutaneous formulations. *Pharmaceutics* 2025; 18(5): 704.
<http://dx.doi.org/10.3390/ph18050704> PMID: 40430523
- [23] Mishra B, Mishra AK, Mohanta YK, et al. Postbiotics: The new horizons of microbial functional bioactive compounds in food preservation and security. *Food Prod Process Nutr* 2024; 6(1): 28.
<http://dx.doi.org/10.1186/s43014-023-00200-w>
- [24] Gupta A, Kumar P. Assessment of the histological state of the healing wound. *Plast Aesthet Res* 2015; 2(5): 239-42.
<http://dx.doi.org/10.4103/2347-9264.158862>
- [25] Karimi F, Montazeri-Najafabady N, Mohammadi F, Azadi A, Koohpeyma F, Gholami A. A potential therapeutic strategy of an innovative probiotic formulation toward topical treatment of diabetic ulcer: An *in vivo* study. *Nutr Diabetes* 2024; 14(1): 66.
<http://dx.doi.org/10.1038/s41387-024-00320-3> PMID: 39164243
- [26] Kim DY, Lee TS, Lee YJ, et al. *Lactobacillus reuteri* NCHBL-005 improves wound healing by promoting the activation of fibroblasts through TLR2/MAPK signaling. *Inflamm Regen* 2025; 45(1): 10.
<http://dx.doi.org/10.1186/s41232-025-00370-9> PMID: 40211423
- [27] Li Z, Zhang S, Zuber F, et al. Topical application of *Lactobacilli* successfully eradicates *Pseudomonas aeruginosa* biofilms and promotes wound healing in chronic wounds. *Microbes Infect* 2023; 25(8): 105176.
<http://dx.doi.org/10.1016/j.micinf.2023.105176> PMID: 37406851
- [28] Mamone V, Fonzo MD, Esposito N, Ferrari M, Ferrari V. Monitoring wound healing with contactless measurements and augmented reality. *IEEE J Transl Eng Health Med* 2020; 8: 1-12.
<http://dx.doi.org/10.1109/JTEHM.2020.2983156> PMID: 32373400
- [29] Santos TS, Santos IDD, Pereira-Filho RN, et al. Histological evidence of wound healing improvement in rats treated with oral administration of hydroalcoholic extract of *Vitis labrusca*. *Curr Issues Mol Biol* 2021; 43(1): 335-52.
<http://dx.doi.org/10.3390/cimb43010028> PMID: 34208147
- [30] Grubbs H, Manna B. Wound physiology. StatPearls. Treasure Island: StatPearls Publishing 2023.
- [31] Wallace HA, Basehore BM, Zito PM. Wound healing phases. StatPearls. Treasure Island: StatPearls Publishing 2023.
- [32] Rosen RD, Manna B. Wound dehiscence. StatPearls. Treasure Island: StatPearls Publishing 2023.
- [33] Wound healing. 2025. Available from: https://www.physiopeia.com/index.php?title=Wound_Healing&ol=1&ol2=1&ol3=1&ol4=1&ol5=1&ol6=1&ol7=1&ol8=1&ol9=1&ol10=1&ol11=1&ol12=1&ol13=1&ol14=1&ol15=1&ol16=1&ol17=1&ol18=1&ol19=1&ol20=1&ol21=1&ol22=1&ol23=1&ol24=1&ol25=1&ol26=1&ol27=1&ol28=1&ol29=1&ol30=1&ol31=1&ol32=1&ol33=1&ol34=1&ol35=1&ol36=1&ol37=1&ol38=1&ol39=1&ol40=1&ol41=1&ol42=1&ol43=1&ol44=1&ol45=1&ol46=1&ol47=1&ol48=1&ol49=1&ol50=1
- [34] Cañedo-Dorantes L, Cañedo-Ayala M. Skin acute wound healing: A comprehensive review. *Int J Inflamm* 2019; 2019: 1-15.
<http://dx.doi.org/10.1155/2019/3706315> PMID: 31275545
- [35] Frykberg RG, Banks J. Challenges in the treatment of chronic wounds. *Adv Wound Care* 2015; 4(9): 560-82.
<http://dx.doi.org/10.1089/wound.2015.0635> PMID: 26339534
- [36] Guo S, DiPietro LA. Factors affecting wound healing. *J Dent Res* 2010; 89(3): 219-29.
<http://dx.doi.org/10.1177/0022034509359125> PMID: 20139336
- [37] Abdulkhaleq LA, Assi MA, Abdullah R, Zamri-Saad M, Taufiq-Yap YH, Hezmee MNM. The crucial roles of inflammatory mediators in inflammation: A review. *Vet World* 2018; 11(5): 627-35.
<http://dx.doi.org/10.14202/vetworld.2018.627-635> PMID: 29915501
- [38] Soliman AM, Barreda DR. Acute inflammation in tissue healing. *Int J Mol Sci* 2022; 24(1): 641.
<http://dx.doi.org/10.3390/ijms24010641> PMID: 36614083
- [39] Wang XJ, Han G, Owens P, Siddiqui Y, Li AG. Role of TGF β -mediated inflammation in cutaneous wound healing. *J Invest Dermatol Symp Proc* 2006; 11(1): 112-7.
<http://dx.doi.org/10.1038/sj.jidsymp.5650004> PMID: 17069018
- [40] Mascharak S, Talbott HE, Januszyk M, et al. Multi-omic analysis reveals divergent molecular events in scarring and regenerative wound healing. *Cell Stem Cell* 2022; 29(2): 315-327.e6.
<http://dx.doi.org/10.1016/j.stem.2021.12.011> PMID: 35077667
- [41] Le Noci V, Bernardo G, Bianchi F, Tagliabue E, Sommariva M, Sfondrini L. Toll like receptors as sensors of the tumor microbial dysbiosis: Implications in cancer progression. *Front Cell Dev Biol* 2021; 9: 732192.
<http://dx.doi.org/10.3389/fcell.2021.732192> PMID: 34604233
- [42] Watanabe S, Zenke K, Muroi M. Lipoteichoic acid inhibits lipopolysaccharide-induced TLR4 signaling by forming an inactive TLR4/MD-2 complex dimer. *J Immunol* 2023; 210(9): 1386-95.
<http://dx.doi.org/10.4049/jimmunol.2200872> PMID: 36897262
- [43] Xiong L, Zhevlakova I, West XZ, et al. TLR2 regulates hair follicle cycle and regeneration via BMP signaling. *eLife* 2024; 12: RP89335.
<http://dx.doi.org/10.7554/eLife.89335.3> PMID: 38483447
- [44] Nguyen MT, Götz F. Lipoproteins of Gram-positive bacteria: Key players in the immune response and virulence. *Microbiol Mol Biol Rev* 2016; 80(3): 891-903.
<http://dx.doi.org/10.1128/MMBR.00028-16> PMID: 27512100
- [45] Guo D, Wang Q, Li C, Wang Y, Chen X. VEGF stimulated the angiogenesis by promoting the mitochondrial functions. *Oncotarget* 2017; 8(44): 77020-7.
<http://dx.doi.org/10.18632/oncotarget.20331> PMID: 29100366
- [46] Xu M, Jiang Z, Wang C, et al. Acetate attenuates inflammasome activation through GPR43-mediated Ca²⁺-dependent NLRP3 ubiquitination. *Exp Mol Med* 2019; 51(7): 1-13.
<http://dx.doi.org/10.1038/s12276-019-0276-5> PMID: 31337751
- [47] Tan BL, Norhaizan ME, Liew WPP, Sulaiman Rahman H. Antioxidant and oxidative stress: A mutual interplay in age-related diseases. *Front Pharmacol* 2018; 9: 1162.
<http://dx.doi.org/10.3389/fphar.2018.01162> PMID: 30405405
- [48] Xie N, Zhang L, Gao W, et al. NAD⁺ metabolism: Pathophysiologic mechanisms and therapeutic potential. *Signal Transduct Target Ther* 2020; 5(1): 227.
<http://dx.doi.org/10.1038/s41392-020-00311-7> PMID: 33028824
- [49] Ukaegbu K, Allen E, Svoboda KKH. Reactive oxygen species and antioxidants in wound healing: Mechanisms and therapeutic potential. *Int Wound J* 2025; 22(5): e70330.

- <http://dx.doi.org/10.1111/iwj.70330> PMID: 40288766
- [50] Lee IT, Lee CW, Tung WH, *et al.* Cooperation of TLR2 with MyD88, PI3K, and Rac1 in lipoteichoic acid-induced cPLA2/COX-2-dependent airway inflammatory responses. *Am J Pathol* 2010; 176(4): 1671-84.
<http://dx.doi.org/10.2353/ajpath.2010.090714> PMID: 20167866
- [51] Lushchak VI. Glutathione homeostasis and functions: Potential targets for medical interventions. *J Amino Acids* 2012; 2012: 1-26.
<http://dx.doi.org/10.1155/2012/736837> PMID: 22500213
- [52] Chai YC, Mieyal JJ. Glutathione and glutaredoxin—Key players in cellular redox homeostasis and signaling. *Antioxidants* 2023; 12(8): 1553.
<http://dx.doi.org/10.3390/antiox12081553> PMID: 37627548
- [53] Forman HJ, Zhang H, Rinna A. Glutathione: Overview of its protective roles, measurement, and biosynthesis. *Mol Aspects Med* 2009; 30(1-2): 1-12.
<http://dx.doi.org/10.1016/j.mam.2008.08.006> PMID: 18796312
- [54] Leser T, Baker A. Molecular mechanisms of *Lactocaseibacillus rhamnosus* LGG® probiotic function. *Microorganisms* 2024; 12(4): 794.
<http://dx.doi.org/10.3390/microorganisms12040794> PMID: 38674738
- [55] von Ossowski I, Reunanen J, Satokari R, *et al.* Mucosal adhesion properties of the probiotic *Lactobacillus rhamnosus* GG SpaCBA and SpaFED pilin subunits. *Appl Environ Microbiol* 2010; 76(7): 2049-57.
<http://dx.doi.org/10.1128/AEM.01958-09> PMID: 20118368
- [56] Widjaja M, Harvey KL, Hagemann L, *et al.* Elongation factor Tu is a multifunctional and processed moonlighting protein. *Sci Rep* 2017; 7(1): 11227.
<http://dx.doi.org/10.1038/s41598-017-10644-z> PMID: 28894125

DISCLAIMER: The above article has been published, as is, ahead-of-print, to provide early visibility but is not the final version. Major publication processes like copyediting, proofing, typesetting and further review are still to be done and may lead to changes in the final published version, if it is eventually published. All legal disclaimers that apply to the final published article also apply to this ahead-of-print version.