



# Estimation of Biofilm Components and Prevalence of SDR Genes among *Staphylococcus aureus* Isolated from Anbar Hospitals

Rafal Abdul Khaliq Juma Almaadhidy<sup>1</sup> and Safaa Abed Lateef Al Meani<sup>2,\*</sup>

<sup>1</sup>Department of Biology, College of Science, University of Anbar, Ramadi 31001, Iraq

<sup>2</sup>Department of Biotechnology, College of Science, University of Anbar, Ramadi 31001, Iraq

© 2024 The Author(s). Published by Bentham Open.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



\*Address correspondence to this author at the Department of Biology, College of Science, University of Anbar, Ramadi 31001, Iraq; E-mail: raf22s1002@uoanbar.edu.iq

Published: November 15, 2024



Send Orders for Reprints to  
[reprints@benthamscience.net](mailto:reprints@benthamscience.net)

**Table 1. Relationship of different genotypes and the ability of biofilm formation (BF) in *Staphylococcus aureus* isolates with individual pulsotypes.**

No. of Strains	Degree of BF	sdrC	Genes	sdrD	sdrE
1	+++	+		+	+
2	+++	+		+	-
3-7	+++	+		+	+
8	+	+		-	-
9	++	+		+	+
10	++	+		-	+
11-12	+++	+		+	+
13	++	+		-	+
14	++	+		+	+
15	++	+		-	+
16_17	+++	+		+	+
18	++	+		+	+
19_22	++	+		-	+
23	+++	+		+	+
24_25	+	+		-	+
26_28	+++	+		+	+
29_30	++	+		+	+
31_32	++	+		-	+
33_35	+++	+		+	+
36_41	++	+		-	+
42	++	+		+	+
43_46	++	+		-	+
47_48	+++	+		+	+

-	-	-	Genes	-
No. of Strains	Degree of BF	<i>sdrC</i>	<i>sdrD</i>	<i>sdrE</i>
49_51	+++	-	+	+
52_53	+	-	-	-
54_55	+	+	-	-
56	++	+	+	+
57_59	++	+	-	+
60_62	+	+	-	-
63_68	++	+	+	-
69_75	++	+	+	+
76_78	++	+	-	-