



# The Open Microbiology Journal

Content list available at: <https://openmicrobiologyjournal.com>



## RESEARCH ARTICLE

### Identification of Novel Mobile Genetic Elements Associated with Resistance to Macrolide and Lincosamide in *Streptococcus dysgalactiae* subsp. *equisimilis*

Alexandra Kireeva<sup>1</sup> and Alexander Dmitriev<sup>1,\*</sup>

<sup>1</sup>Department of Molecular Microbiology, Institute of Experimental Medicine, Saint-Petersburg, Russia

#### Abstract:

#### Background:

*Streptococcus dysgalactiae* subsp. *equisimilis* (SDSE) is an important human pathogen. Recently, several studies have described the incidence of antibiotic resistance for SDSE worldwide, however, the data on the presence of corresponding genes and their possible association with mobile genetic elements are still limited.

#### Objective:

The objective of this research was to analyze the macrolide resistance in SDSE and to identify genetic determinants, mechanisms of resistance, and association with mobile genetic elements.

#### Methods:

A total of 9 SDSE strains from the collection of Joint Russian-Vietnamese Tropical Research and Technological Center (Hanoi, Vietnam) were used. These strains were previously isolated from throat swabs of children with pharyngotonsillitis in 6 provinces in Vietnam from 2012 to 2015. Antimicrobial resistance was tested by disk diffusion method. The presence of antibiotic resistance genes (ARG) was analyzed by PCR. The strains were characterized by *emm* typing and multilocus sequence typing (MLST). Illumina sequencing was employed for genome analysis of 4 representative SDSE isolates. Analysis of genetic elements with antibiotic resistance determinants was done using PubMed database and BLAST-searches. Artemis was used for comparative analysis of genetic elements.

#### Results:

In our study, we identified *emm* types that were similar to those reported in other studies. All SDSE isolates remained susceptible to penicillin, but presented alarming level of resistance to macrolides, tetracyclines, and fluoroquinolones. Most of the erythromycin-resistant strains were also characterized by clindamycin-resistance (MLSB phenotype). Both *erm* and different alleles of *mef* genes widely distributed among *streptococcus pyogenes* and *Streptococcus pneumoniae* were detected, except *erm* (TR) gene. The genetic elements carrying resistance determinants showed significant interspecies similarities, indicating conjugative transfer of antibiotic resistance genes between streptococcal species.

#### Conclusion:

Identification of the novel antibiotic resistance genes in SDSE indicates the necessity of monitoring of antibiotic resistance spreading and gene transfer in this bacterium.

**Keywords:** *Streptococcus dysgalactiae* subsp. *equisimilis*, Macrolide resistance, Clindamycin resistance, Resistance determinants, Mobile genetic elements, Molecular typing, Whole genome sequencing.

#### Article History

Received: August 26, 2022

Revised: December 13, 2022

Accepted: December 15, 2022

## 1. INTRODUCTION

The human  $\beta$ -hemolytic *streptococci* (BHS) include *streptococcus pyogenes* (group A *streptococci*, GAS),

*streptococcus agalactiae* (group B *streptococci*, GBS), and *Streptococcus dysgalactiae* subsp. *equisimilis* (group C and G *streptococci*, SDSE). SDSE colonizes the skin or mucosal surfaces, such as the respiratory tract, gastrointestinal tract, or vagina and causes a variety of diseases including invasive diseases. During the recent years the incidence of invasive SDSE diseases has increased, and in some geographic regions

\* Address correspondence to this author at the Department of Molecular Microbiology, Institute of Experimental Medicine, Saint-Petersburg, Russia; Tel: +7-950-0336655; E-mail: [admitriev10@yandex.ru](mailto:admitriev10@yandex.ru)

the rate of SDSE diseases was higher than those of GAS and GBS [1 - 3].

At present, penicillin is considered as drug of choice for treatment of  $\beta$ -hemolytic streptococcal (BHS) infections. Additionally, macrolides and clindamycin can be used for treatment of the patients intolerant to  $\beta$ -lactam antibiotics. Importantly, combined antibiotic therapy can reduce mortality in case of severe disease manifestations [4], probably through the abrogation of toxin synthesis [5]. Significant variations in resistance to macrolides, lincosamides (such as clindamycin) and streptogramin B (MLSB) in BHS were reported. MLSB resistance in  $\beta$ -hemolytic *streptococci* is mostly provided by the target modification enzymes encoded by *erm* genes, resulting in the resistance to all three classes (MLSB), or *mef* genes encoding efflux pumps specific for macrolides [6]. Lincosamide resistance genes *lsa* and *lnu* also occur in *streptococci* [7]. In GAS and GBS, all these resistance genes are located within mobile genetic elements. Their spreading among *streptococci* occurs by either horizontal genetic transfer or clonal expansion [6 - 9], however, their dissemination has not been extensively studied. *In vitro* studies showed the existence of conjugal transfer of integrative conjugative elements (ICEs) harboring resistance genes between streptococcal species [8].

As a consequence, the goal of the present study was to analyze the current status of antimicrobial resistance in SDSE. In addition, genome sequencing was performed to identify antibiotic resistance genes and their possible association with mobile genetic elements.

## 2. MATERIALS AND METHODS

### 2.1. Bacterial Isolates

A total of 9 SDSE strains from the collection of Joint Russian-Vietnamese Tropical Research and Technological Center (Hanoi, Vietnam) were used. These strains were previously isolated during 2012-2015 from 1359 children of 7-10 years old from different regions of Vietnam [10]. Bacteria were grown on Columbia base agar with 5% of sheep blood and in Todd-Hewitt broth with 5% of inactivated horse serum in 5% CO<sub>2</sub> atmosphere at 37°C. Bacterial DNA was isolated by phenol/chloroform extraction. Species identity was identified with 16S rRNA gene sequencing [11].

### 2.2. *emm* Typing

*emm* typing of SDSE isolates was performed in accordance with recommendations available at CDC web site ([www.cdc.gov/streplab/groupa-strep/emm-typing-protocol.html](http://www.cdc.gov/streplab/groupa-strep/emm-typing-protocol.html)).

### 2.3. Antimicrobial Susceptibility Testing

The isolates were tested for susceptibility to penicillin G, cefotaxime, vancomycin, amikacin, norfloxacin, erythromycin, clindamycin, and tetracycline by disk diffusion method according to EUCAST guidelines. MLSB resistance phenotype was analyzed by double disc diffusion method [12]. Depending on resistance to erythromycin and/or clindamycin, the SDSE isolates were divided into several groups/phenotypes:

constitutive MLSB-resistance (cMLSB), inducible MLSB-resistance (iMLSB), macrolide resistance (M-phenotype), and lincosamide resistance (L phenotype). *Streptococcus pneumoniae* strain ATCC 49619 was used as a control.

### 2.4. Antibiotic Resistance Gene Detection

PCR detection of antibiotic resistance genes *erm*(B), *erm*(TR), *mef*(A/E) (resistance to macrolides, and *tet*(M), *tet*(O), *tet*(T), *tet*(S) (resistance to tetracycline) was done using the primers previously published (Table S1).

### 2.5. Bioinformatic Analysis

The genome sequencing was performed for NT15, V123, B82 isolates with reduced susceptibility to erythromycin or clindamycin, and T201 isolate susceptible to erythromycin and clindamycin. Construction of the libraries and DNA sequencing on MiSeq platform was done as recommended by the manufacturer (Illumina, Essex, United Kingdom). Quality of the reads was tested using FastQC, trimmed with Trimmomatic [13], assembled by Spades [14], and subsequently annotated by the NCBI Prokaryotic Genome Annotation Pipeline (PGAP).

Multilocus sequence typing (MLST) was done using the Center for Genomic Epidemiology website [15]. Relevant resistance genes were identified using the ResFinder and CARD databases [16, 17]. Verification of potential integrative conjugative elements (ICEs) was performed using ICEberg 2.0 [18] together with PubMed database and BLAST-searches. Artemis was used for comparative analysis of genetic elements [19].

### 2.6. Nucleotide Sequence Accession Numbers

The Whole Genome Shotgun (WGS) projects reported in this paper have been deposited at DDBJ/ENA/GenBank (JAFELD000000000 (V123), JAFELE000000000 (B82), JAFELF000000000 (NT15), and JAFELG000000000 (T201)).

## 3. RESULTS

As previously published, a total of 152  $\beta$ -hemolytic *streptococci* were isolated from 1359 children of 7-10 years old from different regions of Vietnam in the period 2012-2015 [10]. GAS were isolated from 49 of 1359 (3,6%) examined children, while group C and G *streptococci* were isolated from 8 (0,6%) and 75 (5,5%) children, respectively [10]. Using *cpn60* gene based PCR approach [20] for differentiation of the species within groups C and G, a total of 9 SDSE strains (1 – group C, and 8 – group G) were identified. Other identified strains included group C *S. anginosus* (4 isolates), *S. parasanguinis* (1 isolate), *S. constellatus* (1 isolate), *S. gardonii* (1 isolate), and group G *S. anginosus* (54 isolates), *S. constellatus* (4 isolates), *S. parasanguinis* (3 isolates), *S. sanguinis* (3 isolates), *S. mitis* (2 isolates), *S. australis* (1 isolate).

### 3.1. SDSE Genetic Diversity

A high clonal diversity was discovered: a total of 6 *emm* types were identified among the 9 SDSE isolates, and *stC5345 emm* type was specific for 3 out of 9 isolates. MLST performed for 4 sequenced SDSE isolates revealed 3 different previously published STs (Table 1).

**Table 1. Antimicrobial resistance phenotypes and genotypes of SDSE isolates.**

Isolate SDSE	<i>emm</i>	MLST	Phenotype	Genotype	ICE-family	Other Resistance Genes
B159	<i>stC5345</i>	nt	MLSB sensitive	-	-	<i>tet(M)</i>
HF196		nt	MLSB sensitive	-	-	<i>tet(M)</i>
<b>T201</b>		44	MLSB sensitive	-	-	<i>tet(M)</i> , <i>pat(B)</i>
HF112	<i>stG480</i>	nt	cMLSB	<i>erm(B)</i>	Tn6002	<i>tet(M)</i>
<b>NT15</b>		323	cMLSB	<i>mef(G)</i>	ΦNT15	<i>tet(S)</i> , <i>lnu(B)</i> , <i>lsa(E)</i> , <i>pat(B)</i>
<b>V123</b>	<i>stG6</i>	44	M	<i>mef(A)</i>	Φ46.1-like	<i>tet(O)</i> , <i>lnu(C)</i> , <i>pat(B)</i>
V63	<i>emm44</i>	nt	cMLSB	<i>erm(B)</i>	Tn917	
T122	<i>stG4831</i>	nt	M	<i>mef(E)</i>	mega	
<b>B82</b>	<i>stC36</i>	499	cMLSB	<i>erm(B)</i>	ICE-B82	<i>tet(T)</i> , <i>pat(B)</i>

Note: Highlighted strains were chosen for whole genome sequencing. ICE, integrative conjugative element; MLST, multilocus sequence typing; MLSB, macrolide, lincosamide, streptogramin B; cMLSB, constitutive MLSB -resistance, M, macrolide resistance alone; mega, macrolide efflux genetic assembly element containing; nt – not tested.

### 3.2. Antimicrobial Susceptibility

All SDSE isolates of this study were susceptible to cefotaxime, vancomycin, penicillin G and resistant to amikacin. The incidence of fluoroquinolones resistance (norfloxacin) was detected in 4 isolates that corresponded with the presence of *pat(B)* gene (Table 1). Resistance to tetracycline was observed in 6 isolates. The overall prevalence of resistance to MLSB antibiotics among SDSE was very high (6 isolates were erythromycin resistant, and 4 of them – additionally resistant to clindamycin (cMLS phenotype)).

### 3.3. Detection of Antibiotic Resistance Genes

In 6 SDSE isolates with M or cMLSB phenotypes (reduced susceptibility to MLSB) the corresponding resistance genes were detected (Table 1). Almost all isolates (3 out of 4) with cMLSB phenotype possessed the *erm(B)* gene, while *mef(G)* was discovered in one isolate which additionally possessed clindamycin resistance genes *lsa(E)* and *lnu(B)*.

The M phenotype specific for 2 isolates was associated with *mef(E)* and *mef(A)* genes, and clindamycin resistance gene *lnu(C)* was detected in *mef(A)* positive isolate.

Among the 9 SDSE isolates, 4 isolates demonstrated co-resistance to tetracycline. *tet(M)* was the most common, while *tet(O)*, *tet(S)* and *tet(T)* were also discovered (each by one) (Table 1).

Furthermore, we used whole genome sequencing of 4 SDSE isolates to identify mobile elements responsible for the spreading of antibiotic resistance genes. In these isolates, the gene *pat(B)* encoding ATP-binding cassette (ABC) for fluoroquinolone antibiotics efflux pump was also discovered.

### 3.4. Mobile Elements Involved in Macrolide and Lincosamide Resistance


#### 3.4.1. Erythromycin-resistant SDSE Isolates Carrying the *mef* Genes

We identified 3 isolates harboring different alleles of *mef* gene (*mefA/E/G*). As expected, in tetracycline-susceptible


isolate T122 PCR assay demonstrated that *mef(E)* gene was located within mega (macrolide efflux genetic assembly) element, which has been previously described for BHS and *Streptococcus salivarius* [21, 22].

In SDSE strain NT15 (GenBank accession number JAFELF000000000) the *mef(G)* gene was located within the prophage-associated genetic element (54,700 bp). This genetic element consisted of 54 open reading frames (ORFs) and was chimeric in nature; it appeared due to the insertion of a Tn1207.1-related transposon into streptococcal prophage. BLASTN analysis revealed the high similarity of this prophage to other prophages, especially Tn1207.3 (52,491 bp) or Φ10394.4 (58,761 bp) discovered in tetracycline-susceptible GAS isolates [23, 24]. In both GAS isolates and SDSE NT15 strain these phages were integrated into the same chromosomal gene, *comEC* (Fig. 1). A Φ10394.4-like element in NT15, entitled ΦNT15, harbored *mef(G)* instead of *mef(A)* and the additional 5 kb region containing unique *lsa(E)* and *lnu(B)* genes. The *lnu(B)* gene is involved in lincosamide modification/inactivation; and the *lsa(E)* encoding ABC transporter is responsible for active efflux of lincosamides, streptogramins A, and pleuromutilins. BLASTN analysis revealed that this region in SDSE strain NT15 is similar to the *lnu(B)*-containing sequences of *S. agalactiae* (JQ861959), *Staphylococcus aureus* (JX560992), *Enterococcus faecalis* (AF408195) and swine *Enterococcus faecium* isolate (KF421157.1) [25]. In addition, in NT15 isolate tetracycline susceptibility is associated with the presence of silent *tet(S)* gene.

The complete sequence of SDSE V123 isolate (GenBank accession number JAFELD000000000) revealed that tetracycline resistance is provided by *tet(O)* gene, and *tet(O)* was linked with *mef(A)* within phage-like element. It is highly similar to well-known Φ46.1 of *S. pyogenes*, which resulted from the insertion of transposon Tn1207.1 into a prophage [26]. However, an addition of IS1595 element and gene *lnu(C)*, which confer resistance to lincosamides, was found (Fig. 2). The Φ46.1-like phage was found to be integrated into a 23S rRNA uracil methyltransferase gene.



**Fig. (1).** An alignment of the phage  $\Phi$ 10394.4 in GAS isolate and the phage  $\Phi$ NT15 in SDSE isolate. The genetic organization  $\Phi$ NT15 was consistent with that of  $\Phi$ 10394.4, a composite element resulting from the insertion of the *mef*-containing Tn1207.1 element into streptococcal prophage. Black arrows, chromosome; purple arrows, ARG (antimicrobial resistance genes); blue arrows, prophage-like region.



**Fig. (2).** Schematic representation of the fragment of phage  $\Phi$ 46.1-like in SDSE isolate V123. Purple arrows, ARG; blue arrows, prophage-like region.

### 3.4.2. Erythromycin-resistant SDSE Isolates Carrying *erm(B)* Gene

Among the 6 macrolide resistant isolates, 3 isolates had the *erm(B)* gene and exhibited constitutive resistant phenotype. According to the results of PCR analysis, the *erm(B)* was carried by the Tn917 in isolate V63 [27]. In another *erm(B)*-positive isolate (HF112), which was also resistant to tetracycline, *erm(B)* was associated with *tet(M)* on Tn6002 [28]. The third *erm(B)*-positive isolate, B82, was chosen for genome sequencing. The following analysis of B82 (GenBank accession number JAFELE000000000) revealed the presence of a 45.4-kb element. At present, this element was not found in other SDSE strains, but it has a partial similarity with SDSE strain WCHSDSE-1, which caused the streptococcal outbreak in China in 2013 [29]. Similar element, which contains the *tet(M)* gene instead of *erm(B)*, is present in *Filifactor alocis* ATCC35896. This bacterium can cause periodontal diseases. In place of the 1.5-kb fragment containing *erm(B)* in SDSE, the strain *F. alocis* has a 14.1-kb fragment with *tet(M)* gene.

The 45.4-kb element of isolate B82 appears to be a conjugative transposon because it contains some specific genes, i.e., site-specific recombinase gene, relaxase-encoding gene (*nicK*), an origin of transfer (*oriT*), conjugal transfer coupling protein gene.


Transposition of this element is characterized by the presence of 3-bp direct repeat suggesting that the putative transposons were truly transposable elements (Fig. 3). This transposon was inserted between the genes encoding hypothetical proteins (SDSE167\_0576 and SDSE167\_0577 in strain 167). Additionally, Tn916 was found with gene *tet(T)* instead of *tet(M)* gene.

## 4. DISCUSSION

Based on the genome comparison, the SDSE which belong to group C and G *streptococci*, are closely related to GAS [2, 30, 31]. Almost all SDSE used in this study belonged to GGS. It corresponds to results of the previous studies, wherein Lancefield group G was found to be the most predominant among the human-recovered SDSE [32 - 34].

The *emm* genotyping was successfully performed for all 9 SDSE isolates, and 6 *emm* types were discovered. Four of them (*stG6*, *stG480*, *stC5345*, *stC36*) have already been found in SDSE isolated in North America, Europe and Australia [35], reflecting the successful dissemination of certain *emm* types in human.

Previously the certain correlation between *stG480* and *stG6* types and invasive infections was demonstrated [36]. However, in our study *stG480* and *stG6* strains were non-invasive.



**Fig. (3).** An alignment of the conjugative transposon-like elements in SDSE isolates B82, WCHSDSE-1 and *F. alocis* ATCC 35896. *traE*, conjugal coupling protein; *traC*, type-IV secretion system protein; *iap*, endopeptidase p60 precursor; *topB*, DNA topoisomerase 3; *helicase gene*, helicase/DNA methylase; *rlx*, relaxase/mobilisation nuclease domain protein; *irtA*, iron import ATP-binding/permease protein; *msbA*, putative ABC transporter ATP-binding protein; *ecfT*, energy-coupling factor transporter transmembrane protein; *ykoD*, putative HMP/thiamine import ATP-binding protein; *fisK*, DNA translocase; *nikK*, relaxase; *ermB*, macrolide resistance; *rec*, recombinase; *tcpC*, conjugative transposon protein; *tetM*, tetracycline resistance; *int*, integrase. Purple arrow, *tetM* gene; red arrow, *ermB* gene.

Results of this study demonstrated that beta-lactams are useful for treatment of SDSE infections that correlates with other studies [3, 4, 37]. However, during the recent years the MIC for penicillin was slightly increased for some GAS and SDSE (0.12 and of 0.25 µg/ml), respectively [38]. Tetracycline resistance was noted in 6 of 9 (67%) isolates. The macrolide and lincosamide resistance in SDSE were high, 67% and 45%, respectively. Inducible clindamycin-resistant phenotype (iMLS<sub>B</sub>) was not found among erythromycin-resistant SDSE. Macrolide resistance in SDSE occurs world-wide, e.g., in Hong Kong (24%), in the USA (19%), in Europe (16%) [12, 38]. Furthermore, the percentage of clindamycin-resistant SDSE is significantly higher than in GAS. It should be taken into account during the treatment of streptococcal toxic shock syndrome caused by SDSE. In case of tonsillopharyngitis, fluoroquinolones can be considered as a second choice because the prevalence of fluoroquinolone-resistance among β-hemolytic *streptococci* is still low (1%), and just a few publications reported fluoroquinolone-resistance in SDSE and GAS. Resistance rate to tetracycline is higher than 60%, and for this reason it can no longer be used for empirical treatment of SDSE infections [38].

The resistance genes of macrolides, lincosamides, tetracyclines, and fluoroquinolones were examined in this study. As result, the presence of *erm(A)* (0%), *erm(B)* (33%), *mef(A/E/G)* (33%) differed from those of human-recovered SDSE in China: *erm(A)* (0%), *erm(B)* (78.6%), *mef(A/E)* (5.4%), Korea: *erm(A)* (4.3%), *erm(B)* (20.3%), *mef(A)* (8.7%), and Japan: *erm(A)* (15.5%), *erm(B)* (11.3%), *mef(A)* (2.8%) [39, 40]. Among the tetracycline resistance genes, *tet(M)* was the most common in this study, but *tet(O)*, *tet(S)* and *tet(T)* were discovered (each by one). Other publications confirmed that *tet(M)* was more predominant than *tet(O)*, e.g., in China and Korea (*tet(M)*: 73.2% and 29.0%, and *tet(O)*:

5.4% and 1.4%, respectively) in human SDSE [12, 39, 40]. Mutations in the quinolone resistance determining regions (QRDR) of *gyr(A)* or *par(C)* are considered as a major mechanism of fluoroquinolone resistance [12]. In this study, resistance to fluoroquinolone antibiotics in 4 isolates was associated with the presence gene *pat(B)* mediating antibiotic efflux pump.

The association of macrolide resistant determinants with mobile genetic elements was previously demonstrated for major streptococcal pathogens that explains spreading of antibiotic resistance. The resistance phenotype has been reported to be transferable by conjugation [8, 41] and transduction [42, 43]. In our study most of erythromycin-resistant strains were additionally resistant to clindamycin (phenotype MLS<sub>B</sub>). In SDSE strains we detected a number of novel genetic elements, including the new Φ10394.4-like element characterized by the insertion of *lnu(B)/lsa(E)*-containing sequence; and the new phi-m46.1-like prophage with insertion of *lnu(C)*-containing sequence, which seems to be acquired from other bacterial species [44]. The novel putative transposon carrying *erm(B)* gene was identified in SDSE, which is likely to be acquired from the bacteria of oral microflora. It represents one more example of genetic exchange of antibiotic resistance between gram positive cocci.

**CONCLUSION**

Identification of the novel antibiotic resistance genes in SDSE indicates the necessity of monitoring of antibiotic resistance spreading and gene transfer in this bacterium.

**LIST OF ABBREVIATIONS**

SDSE = *Streptococcus dysgalactiae* subsp. *equisimilis*

MLST = multilocus sequence typing

## ETHICS APPROVAL AND CONSENT TO PARTICIPATE

Not applicable.

## HUMAN AND ANIMAL RIGHTS

No animals and humans were used for studies that are the basis of this research.

## CONSENT FOR PUBLICATION

Not applicable.

## AVAILABILITY OF DATA AND MATERIALS

The data that support the findings of this study are available from the corresponding author, [D.A.V.] on special request.

## FUNDING

This study was done in the frames of projects supported by the Ministry of science and higher education (075-01135-22-00).

## CONFLICT OF INTEREST

Dr. Alexander Dmitriev is on the Editorial Advisory Board of The Open Microbiology Journal.

## ACKNOWLEDGEMENTS

Declared none.

## SUPPLEMENTARY MATERIALS

Supplementary material is available on the Publisher's website.

## REFERENCES

- [1] Bramhachari PV, Kaul SY, McMillan DJ, Shaila MS, Karmarkar MG, Sriprakash KS. Disease burden due to *Streptococcus dysgalactiae* subsp. *equisimilis* (group G and C streptococcus) is higher than that due to *streptococcus pyogenes* among Mumbai school children. *J Med Microbiol* 2010; 59(2): 220-3. [http://dx.doi.org/10.1099/jmm.0.015644-0] [PMID: 19833781]
- [2] Oppegaard O, Mylvaganam H, Kittang BR. Beta-haemolytic group A, C and G streptococcal infections in Western Norway: A 15-year retrospective survey. *Clin Microbiol Infect* 2015; 21(2): 171-8. [http://dx.doi.org/10.1016/j.cmi.2014.08.019] [PMID: 25658557]
- [3] Wajima T, Morozumi M, Hanada S, *et al.* Molecular characterization of invasive *Streptococcus dysgalactiae* subsp. *equisimilis*, Japan. *Emerg Infect Dis* 2016; 22(2): 247-54. [http://dx.doi.org/10.3201/eid2202.141732] [PMID: 26760778]
- [4] Linnér A, Darenberg J, Sjölin J, Henriques-Normark B, Norrby-Teglund A. Clinical efficacy of polyspecific intravenous immunoglobulin therapy in patients with streptococcal toxic shock syndrome: A comparative observational study. *Clin Infect Dis* 2014; 59(6): 851-7. [http://dx.doi.org/10.1093/cid/ciu449] [PMID: 24928291]
- [5] Mascini EM, Jansze M, Schouls LM, Verhoef J, Van Dijk H. Penicillin and clindamycin differentially inhibit the production of pyrogenic exotoxins A and B by group A *streptococci*. *Int J Antimicrob Agents* 2001; 18(4): 395-8. [http://dx.doi.org/10.1016/S0924-8579(01)00413-7] [PMID: 11691576]
- [6] Varaldo PE, Montanari MP, Giovanetti E. Genetic elements responsible for erythromycin resistance in *streptococci*. *Antimicrob Agents Chemother* 2009; 53(2): 343-53. [http://dx.doi.org/10.1128/AAC.00781-08] [PMID: 19001115]
- [7] Hawkins PA, Law CS, Metcalf BJ, *et al.* Cross-resistance to lincosamides, streptogramins A and pleuromutilins in *streptococcus agalactiae* isolates from the USA. *J Antimicrob Chemother* 2017; 72(7): 1886-92. [http://dx.doi.org/10.1093/jac/dkx077] [PMID: 28333320]
- [8] Palmieri C, Magi G, Creti R, *et al.* Interspecies mobilization of an erm(T)-carrying plasmid of *Streptococcus dysgalactiae* subsp. *equisimilis* by a coresident ICE of the ICESa2603 family. *J Antimicrob Chemother* 2013; 68(1): 23-6. [http://dx.doi.org/10.1093/jac/dks352] [PMID: 22949621]
- [9] Zhou W, Yao K, Zhang G, *et al.* Mechanism for transfer of transposon Tn2010 carrying macrolide resistance genes in *Streptococcus pneumoniae* and its effects on genome evolution. *J Antimicrob Chemother* 2014; 69(6): 1470-3. [http://dx.doi.org/10.1093/jac/dku019] [PMID: 24532683]
- [10] Dmitriev AV, Nosik AG, Linh PK, Loan VT, Giang VH, Il'yasov YY. Molecular analysis of group A, C and G *streptococci* isolated from Vietnamese children. Abstracts of XIX Lancefield International Symposium on streptococci and Streptococcal Diseases. Buenos Aires, Argentina. 2014; p. 91.
- [11] Munson MA, Pitt-Ford T, Chong B, Weightman A, Wade WG. Molecular and cultural analysis of the microflora associated with endodontic infections. *J Dent Res* 2002; 81(11): 761-6. [http://dx.doi.org/10.1177/0810761] [PMID: 12407091]
- [12] Oppegaard O, Skrede S, Mylvaganam H, Kittang BR. Emerging threat of antimicrobial resistance in beta-hemolytic *streptococci*. *Frontiers in Microbiology* 2020; 11:797. [http://dx.doi.org/10.3389/fmicb.2020.0079713]
- [13] Nurk S, Lohse M, Antipov D, Usadel B. Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics* 2014; 30(15): 2114-20. [http://dx.doi.org/10.1093/bioinformatics/btu170]
- [14] Nurk S, Bankevich A, Antipov D, *et al.* Assembling single-cell genomes and mini-metagenomes from chimeric MDA products. *J Comput Biol* 2013; 20(10): 714-37. [http://dx.doi.org/10.1089/cmb.2013.0084] [PMID: 24093227]
- [15] Larsen MV, Cosentino S, Rasmussen S, *et al.* Multilocus sequence typing of total-genome-sequenced bacteria. *J Clin Microbiol* 2012; 50(4): 1355-61. [http://dx.doi.org/10.1128/JCM.06094-11] [PMID: 22238442]
- [16] Zankari E, Hasman H, Cosentino S, *et al.* Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 2012; 67(11): 2640-4. [http://dx.doi.org/10.1093/jac/dks261] [PMID: 22782487]
- [17] Alcock BP, Raphenya AR, Lau TTY, *et al.* CARD 2020: Antibiotic resistance surveillance with the comprehensive antibiotic resistance database. *Nucleic Acids Res* 2019; 48(D1): gkz935. [http://dx.doi.org/10.1093/nar/gkz935] [PMID: 31665441]
- [18] Liu M, Li X, Xie Y, *et al.* ICEberg 2.0: An updated database of bacterial integrative and conjugative elements. *Nucleic Acids Res* 2019; 47(D1): D660-5. [http://dx.doi.org/10.1093/nar/gky1123] [PMID: 30407568]
- [19] Carver T, Harris SR, Berriman M. Artemis: An integrated platform for visualization and analysis of high-throughput sequence-based experimental data. *Parkhill J and McQuillan JABioinformatics* 2011; 28(4): 464-9. [http://dx.doi.org/10.1093/bioinformatics/btr703] [PMID: 22199388]
- [20] Il'iasov I, Biswas I, Totolian AA, Dmitriev AV. A method for differential identification of group C and G *streptococci* with PCR. *Klin Lab Diagn* 2011; 2: 40-3. [PMID: 21506385]
- [21] Stadler C, Teuber M. The macrolide efflux genetic assembly of *Streptococcus pneumoniae* is present in erythromycin-resistant *Streptococcus salivarius*. *Antimicrob Agents Chemother* 2002; 46(11): 3690-1. [http://dx.doi.org/10.1128/AAC.46.11.3690-3691.2002] [PMID: 12384396]
- [22] Del Grosso M, Camilli R, Iannelli F, Pozzi G, Pantosti A. The mef(E)-carrying genetic element (mega) of *Streptococcus pneumoniae*: Insertion sites and association with other genetic elements. *Antimicrob Agents Chemother* 2006; 50(10): 3361-6. [http://dx.doi.org/10.1128/AAC.00277-06] [PMID: 17005818]
- [23] Banks DJ, Porcella SF, Barbian KD, Martin JM, Musser JM. Structure and distribution of an unusual chimeric genetic element encoding macrolide resistance in phylogenetically diverse clones of group A

- Streptococcus. J Infect Dis 2003; 188(12): 1898-908. [http://dx.doi.org/10.1086/379897] [PMID: 14673771]
- [24] Giovanetti E, Brenciani A, Vecchi M, Manzin A, Varaldo PE. Prophage association of *mef(A)* elements encoding efflux-mediated erythromycin resistance in *streptococcus pyogenes*. J Antimicrob Chemother 2005; 55(4): 445-51. [http://dx.doi.org/10.1093/jac/dki049] [PMID: 15772148]
- [25] Montilla A, Zavala A, Cáceres Cáceres R, et al. Genetic environment of the *lnu(B)* gene in a *streptococcus agalactiae* clinical isolate. Antimicrob Agents Chemother 2014; 58(9): 5636-7. [http://dx.doi.org/10.1128/AAC.02630-14] [PMID: 24957835]
- [26] Brenciani A, Bacciaglia A, Vignaroli C, Pugnali A, Varaldo PE, Giovanetti E. *Phim46.1*, the main *streptococcus pyogenes* element carrying *mef(A)* and *tet(O)* genes. Antimicrob Agents Chemother 2010; 54(1): 221-9. [http://dx.doi.org/10.1128/AAC.00499-09] [PMID: 19858262]
- [27] Shaw JH, Clewell DB. Complete nucleotide sequence of macrolide-lincosamide-streptogramin B-resistance transposon Tn917 in *Streptococcus faecalis*. J Bacteriol 1985; 164(2): 782-96. [http://dx.doi.org/10.1128/jb.164.2.782-796.1985] [PMID: 2997130]
- [28] Warburton PJ, Palmer RM, Munson MA, Wade WG. Demonstration of *in vivo* transfer of doxycycline resistance mediated by a novel transposon. J Antimicrob Chemother 2007; 60(5): 973-80. [http://dx.doi.org/10.1093/jac/dkm331] [PMID: 17855723]
- [29] Wang XH, Zhang XX, Zong ZY. Genome sequence and virulence factors of a group G *Streptococcus dysgalactiae* subsp. *equisimilis* strain with a new element carrying *erm(B)*. Sci Rep 2016; 6(6): 20389. [http://dx.doi.org/10.1038/srep20389]
- [30] Carapetis JR, Steer AC, Mulholland EK, Weber M. The global burden of group A streptococcal diseases. Lancet Infect Dis 2005; 5(11): 685-94. [http://dx.doi.org/10.1016/S1473-3099(05)70267-X] [PMID: 16253886]
- [31] Rantala S. *Streptococcus dysgalactiae* subsp. *equisimilis* bacteremia: An emerging infection. Eur J Clin Microbiol Infect Dis 2014; 33(8): 1303-10. [http://dx.doi.org/10.1007/s10096-014-2092-0] [PMID: 24682845]
- [32] Preziuso S, Pinho MD, Attili AR, et al. PCR based differentiation between *Streptococcus dysgalactiae* subsp. *equisimilis* strains isolated from humans and horses. Comp Immunol Microbiol Infect Dis 2014; 37(3): 169-72. [http://dx.doi.org/10.1016/j.cimid.2014.04.001] [PMID: 24813401]
- [33] Ahmad Y, Gertz RE Jr, Li Z, et al. Genetic relationships deduced from *emm* and multilocus sequence typing of invasive *Streptococcus dysgalactiae* subsp. *equisimilis* and *S. canis* recovered from isolates collected in the United States. J Clin Microbiol 2009; 47(7): 2046-54. [http://dx.doi.org/10.1128/JCM.00246-09] [PMID: 19386831]
- [34] Tanaka D, Isobe J, Watahiki M, et al. Genetic features of clinical isolates of *Streptococcus dysgalactiae* subsp. *equisimilis* possessing lancefield's group a antigen. J Clin Microbiol 2008; 46(4): 1526-9. [http://dx.doi.org/10.1128/JCM.02188-07] [PMID: 18305132]
- [35] McMillan DJ, Bessen DE, Pinho M, et al. Population genetics of *Streptococcus dysgalactiae* subsp. *equisimilis* reveals widely dispersed clones and extensive recombination. Plos One 2010; 5(7): e11741. [http://dx.doi.org/10.1371/journal.pone.0011741]
- [36] Jensen A, Kilian M. Delineation of *Streptococcus dysgalactiae*, its subspecies, and its clinical and phylogenetic relationship to *streptococcus pyogenes*. J Clin Microbiol 2012; 50(1): 113-26. [http://dx.doi.org/10.1128/JCM.05900-11] [PMID: 22075580]
- [37] Pinho MD, Erol E, Ribeiro-Gonçalves B, et al. Beta-hemolytic *Streptococcus dysgalactiae* strains isolated from horses are a genetically distinct population within the *Streptococcus dysgalactiae* taxon. Sci Rep 2016; 6(1): 31736. [http://dx.doi.org/10.1038/srep31736] [PMID: 27530432]
- [38] Brown DFJ, Hope R, Livermore DM, et al. Non-susceptibility trends among enterococci and non-pneumococcal *streptococci* from bacteraemias in the UK and Ireland, 2001-06. J Antimicrob Chemother 2008; 62(S2): ii75-85. [http://dx.doi.org/10.1093/jac/dkn354] [PMID: 18819982]
- [39] Kataja J, Seppälä H, Skurnik M, Sarkkinen H, Huovinen P. Different erythromycin resistance mechanisms in group C and group G *streptococci*. Antimicrob Agents Chemother 1998; 42(6): 1493-4. [http://dx.doi.org/10.1128/AAC.42.6.1493] [PMID: 9624500]
- [40] Zheng PX, Chan YC, Chiou CS, Hsieh CL, Chiang-Ni C, Wu JJ. Highly prevalent *emmSTG840.0* and *emmSTC839.0* types of erythromycin non-susceptible group G *Streptococcus* isolated from bacteremia in southern Taiwan. J Microbiol Immunol Infect 2017; 50(6): 831-8. [http://dx.doi.org/10.1016/j.jmii.2016.12.010] [PMID: 28711431]
- [41] Giovanetti E, Brenciani A, Burioni R, Varaldo PE. A novel efflux system in inducibly erythromycin-resistant strains of *streptococcus pyogenes*. Antimicrob Agents Chemother 2002; 46(12): 3750-5. [http://dx.doi.org/10.1128/AAC.46.12.3750-3755.2002] [PMID: 12435672]
- [42] Ubukata K, Konno M, Fujii R. Transduction of drug resistance to tetracycline, chloramphenicol, macrolides, lincomycin and clindamycin with phages induced from *streptococcus pyogenes*. J Antibiot 1975; 28(9): 681-8. [http://dx.doi.org/10.7164/antibiotics.28.681] [PMID: 1102514]
- [43] Hyder SL, Streitfeld MM. Transfer of erythromycin resistance from clinically isolated lysogenic strains of *streptococcus pyogenes* via their endogenous phage. J Infect Dis 1978; 138(3): 281-6. [http://dx.doi.org/10.1093/infdis/138.3.281] [PMID: 359722]
- [44] Gravey F, Galopin S, Grall N, et al. Lincosamide resistance mediated by *lnu(C)* (L phenotype) in a *Streptococcus anginosus* clinical isolate. J Antimicrob Chemother 2013; 68(11): 2464-7. [http://dx.doi.org/10.1093/jac/dkt255] [PMID: 23812683]