

IgG Endopeptidase SeMac does not Inhibit Opsonophagocytosis of *Streptococcus equi* Subspecies *equi* by Horse Polymorphonuclear Leukocytes

Mengyao Liu and Benfang Lei*

Department of Veterinary Molecular Biology, Montana State University, Bozeman, Montana 59717, USA

Abstract: The secreted Mac protein made by group A *Streptococcus* (GAS) inhibits opsonophagocytosis of GAS by human polymorphonuclear leukocytes (PMNs). This protein also has the endopeptidase activity against human immunoglobulin G (IgG), and the Cys94, His262 and Asp284 are critical for the enzymatic activity. The horse pathogen *Streptococcus equi* subspecies *equi* produces a homologue of Mac (SeMac). SeMac was characterized to determine whether SeMac has IgG endopeptidase activity and inhibits opsonophagocytosis of *S. equi* by horse PMNs. The gene was cloned and recombinant SeMac was overexpressed in *Escherichia coli* and purified to homogeneity. Mice with experimental *S. equi* infection and horses with strangles caused by *S. equi* seroconverted to SeMac, indicating that SeMac is produced *in vivo* during infection. SeMac has endopeptidase activity against human IgG. However, the protein just cleaves a small fraction, which may be IgG1 only, of horse IgG. Replacement of Cys102 with Ser or His272 with Ala abolishes the enzymatic activity of SeMac, and the Asp294Ala mutation greatly decreases the enzymatic activity. SeMac does not inhibit opsonophagocytosis of *S. equi* by horse PMNs but opsonophagocytosis of GAS by human PMNs. Thus, SeMac is a cysteine endopeptidase with a limited activity against horse IgG and must have other function.

Keywords: IgG, Endopeptidase SeMac, opsonophagocytosis, polymorphonuclear leukocytes.

INTRODUCTION

Gram-positive bacterium *Streptococcus equi* subspecies *equi* (*S. equi*) causes equine strangles, a highly contagious purulent lymphadenitis and one of the most common infectious diseases in horses [1,2]. The infection initially causes nasal discharge and fever and, then, leads to abscess formation in local lymph nodes, causing enormous pain and respiratory difficulty. There is massive infiltration of polymorphonuclear leukocytes (PMNs) to the infection site. However, *S. equi* effectively evades the horse innate immunity by being resistant to phagocytosis by PMNs. Horses recovered from strangles acquire immunity against *S. equi* reinfection [3]. The immunity is primarily mediated by protective antibodies [4], which opsonize and thus enhance phagocytosis of *S. equi* by horse PMNs.

To survive in hosts, bacterial pathogens have evolved multiple mechanisms to evade host defense. For examples, both *S. equi* and group A *Streptococcus* (GAS) produce the hyaluronic acid capsule and surface protein M protein to contribute to resistance to phagocytosis by PMNs. We found that GAS produces a secreted Mac protein (also known as IdeE), which inhibits opsonophagocytosis of GAS by human PMNs [5]. This protein can cleave the heavy chain of human immunoglobulin G (IgG) using Cys94, His262 and Asp284 as its catalytic triad [6-8]. There are two kinds of Mac

produced by GAS isolates [7], which use different mechanisms to block the interaction between IgG and Fc receptor on the surface of PMNs. The type-1 Mac, such as M1 Mac produced by serotype M1 GAS strains, has high enzymatic activity and low affinity to Fc receptor on the surface of PMNs, while the type-2 Mac can bind to the Fc receptor and has lower enzymatic activity [9]. *S. equi* has a homologue of GAS M1 Mac (designated SeMac). In this study, SeMac was prepared and characterized. The results indicate that SeMac is a cysteine endopeptidase but does not inhibit opsonophagocytosis of *S. equi* by horse PMNs, suggesting that SeMac has function other than evading horse acquired immunity against *S. equi* infection.

MATERIALS AND METHODS

Materials

Purified horse IgG1 and a mixture of horse IgG1 and IgG4 were kindly provided by Dr. Bettina Wagner at Cornell University. *S. equi*-specific mouse sera were obtained from adult female outbred CD-1 Swiss mice (Charles River Laboratories, Wilmington, Mass.) 21 days after they were inoculated subcutaneously with 1×10^7 cfu *S. equi* strain SEM1. Convalescent sera from 3 horses suffering from strangles were obtained 30 days after diagnosis. GAS M1 Mac was prepared as previously described [5].

Bacterial Strains and Growth

Six of 10 *S. equi* strains used were kindly provided by Dr. James Musser at Methodist Hospital, Houston, Texas,

*Address correspondence to this author at the Department of Veterinary Molecular Biology, Montana State University, P.O. Box 173610, Bozeman, Montana 59717, USA; Tel: 406-994-6389; Fax: 406-994-4303; E-mail: blei@montana.edu

and these strains were isolated more than 20 years ago from horses with strangles in the Eastern U.S. (4 strains), Brazil (1 strain), and Finland (1 strain). The other 4 strains were isolated in 2003 from horses with strangles in Livingston (designated strain SEM1) [10], Pony, Great Fall, and Norris in Montana. GAS strain MGAS5005 (serotype M1) has been described [11]. *Escherichia coli* Novablue and BL21(DE3) (Novagen, Madison, Wis.) were used for gene cloning and protein expression, respectively. *S. equi* and GAS strains were routinely grown in Todd-Hewitt broth (Difco Laboratories, Detroit, Mich.) supplemented with 0.2% yeast extract (THY) in 5% CO₂ at 37°C. THY agar and tryptose agar with 5% sheep blood (Becton Dickinson, Cockeysville, Md.) were used as the solid media.

Gene Cloning and Mutagenesis

The gene fragment encoding mature SeMac was cloned from SEM1 with primers 5'-ACCATGGACGA TTACCAAAGGAATGCTAC-3' and 5'-CGAATTCT TAGCTCAGTTTCTGCCATATG-3'. The protein made from this cloned fragment lacks the presumed secretion signal sequence (amino acids 1-34). The PCR product was digested with *NcoI* and *EcoRI* and ligated into pET-21d (Novagen) to yield recombinant plasmid pSEMAC. The cloned gene was sequenced and had identical DNA sequence with the corresponding open reading frame of *S. equi* genome database [12]. Amino acid replacement of Cys102 and His272 or Asp294 of SeMac with serine and alanine, respectively, was achieved by site-directed mutagenesis using the Quick-Change Mutagenesis kit (Stratagene, La Jolla, Calif.). The entire mutated gene was sequenced to confirm the mutations and rule out spurious mutations.

Expression and Purification of Recombinant SeMac

Recombinant SeMac was purified from *E. coli* BL21 (DE3) containing plasmid pSEMAC. Bacteria were grown to optical density at 600 nm of 0.5 in 6 liters of Luria-Bertani broth supplemented with 100 mg of ampicillin per liter at 37°C, and SeMac expression was then induced with 0.5 mM IPTG 6 h. Solutions used in purification were buffered with 10 mM Tris-HCl (pH 8.0). Cell paste was sonicated for 15 min at 4°C in 60 ml of the buffer, and centrifuged at 15,000 g for 10 min. The supernatant obtained was loaded onto a DEAE-Sepharose column (2.5 by 10 cm). The column was washed with 50 ml of Tris-HCl, and SeMac was eluted with 200 ml of 50 mM NaCl. SeMac was identified by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), and peak fractions were pooled. Ammonium sulfate was added to the pool to a concentration of 1.2 M, and the resulting solution was applied to a phenyl-Sepharose column (1.5 x 10 cm). The column was washed with 50 ml of 1.2 M (NH₄)₂SO₄ and eluted with a 100-ml linear gradient of 1.2 to 0.7 M (NH₄)₂SO₄. Fractions containing the protein were pooled. The pooled protein was precipitated with (NH₄)₂SO₄ at 70% of saturation, centrifuged, dialyzed against 3 liters of 10 mM Tris-HCl buffer (pH 8.0) overnight, and loaded onto a DEAE-Sepharose column (1.5 by 10 cm). The protein was eluted with a 100-ml linear gradient of 2 to 8 mM NaCl and pooled. The pooled protein was concentrated by (NH₄)₂SO₄ precipitation and dialyzed against Tris-HCl as described above.

DNA Sequencing

Genomic DNA was isolated with a FastDNA SPIN Kit (Qbiogene, Calsbad, Calif.) according to manufacturer's protocol. A DNA fragment containing *S. equi mac* was amplified by PCR using the genomic DNA samples and primers 5'-GCATCTCTACTATCTCATCAC-3' and 5'-ACAGGCACATTAATGTTTAAAC-3'. Sequences of the PCR products were obtained from both DNA strands with an Applied Biosystem 310 automated sequencer (Applied BioSystems, Inc., Foster City, Calif.).

Anti-SeMac antisera

Four female outbred CD-1 Swiss mice (4-week-old) (Charles River Laboratories) were immunized subcutaneously with 50 µg of recombinant SeMac suspended in 200 µl of saline emulsified in 44 µl of monophosphoryl lipid A-synthetic trehalose dicorynomycolate adjuvant (Corixa, Hamilton, Mont.). The animals were anesthetized by isoflurane inhalation prior to immunization and blood collection. Blood was collected prior to active immunization to prepare control sera. Mice were boosted at weeks 2 and 4 with 50 µg of protein mixed with the adjuvant. Anti-SeMac antiserum was prepared from blood obtained 1 week after the second booster.

Phagocytosis Assay

Phagocytosis assay was performed as described previously [13, 14]. Briefly, *S. equi* SEM1 and GAS bacteria from exponential growth phase in THY were washed with phosphate-buffered saline (PBS) and labeled with 0.75 µg/ml FITC in PBS at 37°C for 20 min. The labeled bacteria were washed and resuspended at 1 x 10⁹ cfu/ml in PBS. Ten µl of the labeled bacteria were mixed with 100 µl of non-immune heparinized horse or human blood with or without 100 µg/ml SeMac and incubated with gentle shaking at 37°C for 5, 10, or 20 min. The samples were immediately processed using an Immunolyse Kit (Beckman Coulter) according to the manufacturer's protocol and analyzed by flow cytometry. The percentage of PMNs with fluorescent bacteria was used as a measure of phagocytosis efficiency.

Other Assays

To assess IgG endopeptidase activity of SeMac, human, mouse or horse IgG (20 µg) was incubated with *E. coli* lysate containing SeMac proteins or 1 µg purified SeMac or GAS M1 Mac in PBS at 37° for 90 min, and the reaction mixture was analyzed by SDS-PAGE. Western immunoblot analysis, which was performed as described previously [15], was used to assess *in vitro* SeMac production and the presence of SeMac-specific antibody in the horse and mouse convalescent sera. Culture supernatant proteins of *S. equi*, which were used to assess the *in vitro* SeMac production, were prepared by the method of Lei *et al.* [16].

RESULTS

S. equi Mac Gene

S. equi genome encodes a Mac homologue SeMac. The protein shares 62.4% identity in amino acid sequence with Mac made by serotype M1 GAS and a putative secretion

Purified recombinant SeMac and GAS Mac were then tested for activity to cleave horse IgG. While both proteins cleaved human IgG efficiently, only a small fraction of horse IgG was cleaved by either protein (Fig. 4B), suggesting that SeMac or GAS Mac cannot digest all subgroups of horse IgG. To test this possibility, available horse IgG1 and a mixture of IgG1 and IgG4 were treated with SeMac or GAS Mac. SeMac or GAS Mac cleaved a small portion of the IgG1/IgG4 mixture, but not IgG4. Therefore, the Mac proteins can cleave IgG1 but not IgG4. Due to the unavailability of other purified subgroup IgG, it is not known whether SeMac cleaves the other IgG subgroups.




Fig. (4). IgG endopeptidase activity of SeMac. (A) IgG-endopeptidase activity and critical amino acid residues for activity. Human IgG was incubated without (none) or with *E. coli* lysate containing wild-type SeMac (wt), SeMac^{Cys102Ser} (C102S), SeMac^{His272Ala}, or SeMac^{Asp294Ala} at room temperature for 60 min and analyzed by SDS-PAGE. (B) Limited cleavage of horse IgG by SeMac and GAS Mac. Horse total IgG (eIgG), IgG4 (eIgG4), IgG1/IgG4 mixture (eIgG1/4), or human IgG was incubated with purified SeMac or GAS M1 Mac and analyzed by SDS-PAGE.

SeMac does not Inhibit Opsonophagocytosis of *S. equi* by Horse PMN

A whole blood based phagocytosis assay [13, 14] was used to test whether SeMac inhibits opsonophagocytosis of *S. equi* by horse PMNs. The bacteria were labeled with FITC, treated with *S. equi*-specific horse convalescent serum in the absence or presence of SeMac, and incubated with horse blood. Percentage of PMNs with phagocytosed *S. equi* was determined by flow cytometry. As expected, percentage of PMNs with phagocytosed *S. equi* with the serum treatment was significantly higher than that for *S. equi* without serum treatment. But the inclusion of SeMac in the assay did not affect opsonophagocytosis of *S. equi* (Fig. 5A), suggest-

ing that SeMac does not inhibit opsonophagocytosis of *S. equi* by horse PMNs.

The inability of SeMac to inhibit *S. equi* opsonophagocytosis by horse PMNs may be due to its inability to cleave the majority of horse IgG. If this is true, SeMac should inhibit opsonophagocytosis of GAS by human PMNs, since SeMac efficiently cleaves human IgG. To test this idea, the phagocytosis assay was repeated using GAS and human blood. SeMac indeed inhibits the opsonophagocytosis of GAS by human PMNs (Fig. 5B). These results suggest that IgG endopeptidase activity of SeMac is critical for inhibition of phagocytosis.




Fig. (5). Effects of SeMac on opsonophagocytosis of *S. equi* and GAS by horse (A) and human (B) PMNs, respectively. FITC-labeled *S. equi* and GAS bacteria (10^7 c.f.u.) were treated without or with horse and human convalescent sera and incubated in the presence or absence of 100 μ g/ml SeMac with 100 μ l nonimmune horse and human blood, respectively, at 37°C for 5, 10 or 20 min. Red blood cells were lysed using an Immunolyse kit, and percentages of PMNs with phagocytosed bacteria determined by flow cytometry are presented.

DISCUSSION

In this study we describe that SeMac is a cysteine endopeptidase with limited activity against horse IgG and is unable to inhibit opsonophagocytosis of *S. equi* by horse PMNs in a whole blood assay. SeMac efficiently cleaves the heavy chain of human IgG. The residues of Cys102 and His272 are essential and Asp294 important for the enzymatic activity of SeMac. Thus, like GAS M1 Mac [7,8], SeMac is a cysteine endopeptidase. Although both SeMac and GAS Mac efficiently cleave human IgG, they cannot cleave the

Table 1. The Amino Acid Sequences of the Lower Hinge Region of Human and Horse IgG Subgroups and Their Cleavability by GAS M1 Mac and SeMac

IgG Subgroup	AA Sequence in the Lower Hinge Region ^a	Cleaved by ^b	
		M1 Mac	SeMac
Human IgG1	HTCPPCPAPELL <u>GG</u> PSV	+	+
Human IgG2	VE.....PVA- <u> </u>	+	+
Human IgG3	PP..R..... <u> </u>	+	+
Human IgG4	PP..S.....F. <u> </u>	+	+
Horse IgG1	SK..KC.....	+	+
Horse IgG2	PQ...YTHSKF.....	ND	ND
Horse IgG3	CE..K.....	ND	ND
Horse IgG4	GGC.T..PEC.SV....	-	-
Horse IgG5	SP..K.....P.....	ND	ND
Horse IgG6	VIKE..CC.KCP.R...	ND	ND
Horse IgG7	CGGC.TC..C.SV....	ND	ND

^aCleavage was prior to the Gly residue underlined and determined by N-terminal amino acid sequencing of the proteolytic products in the reaction catalyzed by GAS M1, Mac and dots indicate identical residues with human IgG1.

^b+, cleaved; -, not cleaved; ND, not determined.

majority of total horse IgG. The peptide fragment, PELLGG, in the lower hinge region is proposed to bind to the active site of Mac [9], and the cleavage occurs between the two Gly residues (Table 1). This lower hinge region is not well conserved in the seven subgroups of horse IgG [17] (Table 1). Cleavable horse IgG1 has PELLGG, while the non-cleavable horse IgG4 has PECLSVG in the same region, suggesting that the amino acid sequences of the lower hinge region are important for cleavability by SeMac and GAS Mac. If this is true, horse IgG3 may be also cleavable by SeMac, while the other five horse IgG subgroups may not be cleaved. These horse IgG antibodies that cannot be cleaved by SeMac thus still can mediate the opsonophagocytosis of *S. equi*.

SeMac and GAS M1 Mac show similar enzymatic specificity (Fig. 4 and Table 1), confirming the previous finding [18]. SeMac can cleave human IgG and inhibit the opsonophagocytosis of GAS by human PMNs, but it has limited enzymatic activity against horse IgG and is unable to inhibit opsonophagocytosis of *S. equi* by horse PMNs. Thus, there is a correlation between the enzymatic activity of SeMac and its ability to inhibit opsonophagocytosis, suggesting that SeMac functions like GAS M1 Mac in the inhibition of opsonophagocytosis of GAS by human PMNs.

Timoney *et al.* recently reported that IdeE/SeMac reduces the bactericidal activity of isolated equine PMNs for *S. equi* [19]. Our results suggest that the inhibition of the bactericidal activity of PMNs may not be mediated by opsonophagocytosis or may be insignificant in whole blood.

SeMac is not produced *in vitro*, whereas GAS Mac is [16]. The *mac* gene is controlled by the two-component regulatory system CovRS [5], which also controls the expression of many virulence factors including the hyaluronic capsule [20]. The hyaluronic capsule of *S. equi* is highly

produced *in vitro*. These observations suggest that the *semac* and *mac* genes are regulated by different mechanisms. This suggestion is supported by only 29% DNA sequence identity between *S. equi* and M1 GAS in the upstream region of the *semac* and *mac* genes.

S. equi is a horse pathogen. The fact that SeMac does not inhibit opsonophagocytosis of *S. equi* by horse PMNs indicates that SeMac is not involved in the evasion of the acquired horse immunity against *S. equi*. This suggests that SeMac has other unknown function.

ACKNOWLEDGEMENTS

This work was supported by the National Research Initiative of the USDA Cooperative State Research, Education and Extension Service, grant number 2007-35204-18306), the National Center for Research Resources, grant P20 RR-020185, and the Montana State University Agricultural Experimental Station and USDA Formula Funds. We thank Drs. James Musser and Bettina Wagner for providing horse *S. equi* strains and purified horse IgG, respectively.

REFERENCES

- [1] Timoney JF. Strangles. *Vet Clin North Am Equine Pract* 1993; 9: 365-74.
- [2] Harrington DJ, Sutcliffe IC, Chanter N. The molecular basis of *Streptococcus equi* infection and disease. *Microbes Infect* 2002; 4: 501-10.
- [3] Woolcock JB. Immunity to *S. equi*. *Aust Vet J* 1975; 51: 554-9.
- [4] Timoney JF, Eggers D. Serum bactericidal responses to *Streptococcus equi* of horses following vaccination. *Equine Vet J* 1985; 17: 306-10.
- [5] Lei B, DeLeo FR, Hoe NP, *et al.* Evasion of human innate and acquired immunity by a bacterial homologue of CD11b that inhibits opsonophagocytosis. *Nat Med* 2001; 7: 1298-305.
- [6] von Pawel-Rammingen U, Johansson BP, Björck L. IdeS, a novel streptococcal cysteine proteinase with unique specificity for immunoglobulin G. *EMBO J* 2002; 21: 1607-15.

- [7] Lei B, DeLeo FR, Reid SD, *et al.* Opsonophagocytosis-inhibiting Mac protein of group A Streptococcus: identification and characteristics of two genetic complexes. *Infect Immun* 2002; 70: 6880-90.
- [8] Lei B, Liu M, Meyers EG, Manning HM, Nagiec MJ, Musser JM. histidine and aspartic acid residues important for IgG endopeptidase activity of the group A *Streptococcus* opsonophagocytosis-inhibiting mac protein. *Infect Immun* 2003; 71: 2881-4.
- [9] Agniswamy J, Lei B, Musser JM, Sun PD. Insight of host immune evasion mediated by two variants of Group A *streptococcus* Mac protein. *J Biol Chem* 2004; 279: 52789-96.
- [10] Liu M, McClure MJ, Zhu H, Xie G, Lei B. The two-component regulatory system VicRK is important to virulence of *Streptococcus equi* subspecies *equi*. *Open Microbiol J* 2008; 2: 89-93.
- [11] Hoe NP, Nakashima K, Grigsby D, *et al.* Rapid molecular genetic subtyping of serotype M1 group A *Streptococcus* strains. *Emerg Infect Dis* 1999; 5:254-263.
- [12] Holden MT, Heather Z, Paillot R, *et al.* Genomic evidence for the evolution of *Streptococcus equi*: host restriction, increased virulence, and genetic exchange with human pathogens. *PLoS Pathog* 2009; 5: e1000346.
- [13] White-Owen C, Alexander JW, Sramkoski RM, Babcock GF. Rapid whole-blood microassay using flow cytometry for measuring neutrophil phagocytosis. *J Clin Microbiol* 1992; 30: 2071-6.
- [14] Liu M, Hanks TS, Zhang J, *et al.* Defects in ex vivo and in vivo growth and sensitivity to osmotic stress of group A *Streptococcus* caused by interruption of response regulator gene *vicR*. *Microbiology* 2006; 152: 967-78.
- [15] Xie G, Liu M, Zhu H, Lei B. Esterase SeE of *Streptococcus equi* ssp. *equi* is a novel non-specific carboxylic ester hydrolase. *FEMS Microbiol Lett* 2008; 289: 181-6.
- [16] Lei B, Mackie S, Lukomski S, Musser JM. Identification and immunogenicity of Group A *Streptococcus* culture supernatant proteins. *Infect Immun* 2000; 68: 6807-18.
- [17] Wagner B. Immunoglobulins and immunoglobulin genes of the horse. *Dev Comp Immunol* 2006; 30:155-64.
- [18] Lannergård J, Guss B. IdeE, an IgG-endopeptidase of *Streptococcus equi* ssp. *equi*. *FEMS Microbiol Lett* 2006; 262: 230-5.
- [19] Timoney JF, Yang J, Liu J, Merant C. IdeE reduces the bactericidal activity of equine neutrophils for *Streptococcus equi*. *Vet Immunol Immunopathol* 2008; 122: 76-82.
- [20] Heath A, DiRita VJ, Barg NL, Engleberg NC. A two-component regulatory system, CsrR-CsrS, represses expression of three *Streptococcus pyogenes* virulence factors, hyaluronic acid capsule, streptolysin S, and pyrogenic exotoxin B. *Infect Immun* 1999; 67: 5298-305.

Received: March 01, 2010

Revised: March 12, 2010

Accepted: March 15, 2010

© Liu and Lei; Licensee *Bentham Open*.

This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.