

Genomic Analysis Reveals Versatile Organisms for Quorum Quenching Enzymes: Acyl-Homoserine Lactone-Acylase and -Lactonase

Vipin Chandra Kalia^{1,*}, Sajan C. Raju² and Hemant J. Purohit²

¹Microbial Biotechnology and Genomics, Institute of Genomics and Integrative Biology (IGIB), CSIR, Delhi University Campus, Mall Road, Delhi-110007, India; ²Environmental Genomics Unit, National Environmental Engineering Research Institute (NEERI), CSIR, Nehru Marg, Nagpur - 440020, India

Abstract: Microbial virulence and their resistance to multiple drugs have obliged researchers to look for novel drug targets. Virulence of pathogenic microbes is regulated by signal molecules such as acylated homoserine lactone (AHL) produced during a cell density dependent phenomenon of quorum sensing (QS). In contrast, certain microbes produce AHL-lactonases and -acylases to degrade QS signals, also termed as quorum quenching. Mining sequenced genome databases has revealed organisms possessing conserved domains for AHL-lactonases and -acylases: i) *Streptomyces* (Actinobacteria), ii) *Deinococcus* (Deinococcus-Thermus), iii) *Hyphomonas* (α -Proteobacteria), iv) *Ralstonia* (β -Proteobacteria), v) *Photorhabdus* (γ -Proteobacteria), and certain marine gamma proteobacterium. Presence of genes for both the enzymes within an organism was observed in the following: i) *Deinococcus radiodurans* R1, ii) *Hyphomonas neptunium* ATCC 15444 and iii) *Photorhabdus luminescens* subsp. *laumondii* TTO1. These observations are supported by the presence motifs for lactonase and acylase in these strains. Phylogenetic analysis and multiple sequence alignment of the gene sequences for AHL-lactonases and -acylases have revealed consensus sequences which can be used to design primers for amplifying these genes even among mixed cultures and metagenomes. Quorum quenching can be exploited to prevent food spoilage, bacterial infections and bioremediation.

Keywords: Acylhomoserine lactone; Acylase; *Bacillus*; Lactonase; Phylogeny; *Pseudomonas*; Quorum sensing; Quorum quenching.

INTRODUCTION

The advent of antibiotics in the beginning of twentieth century gave a new lease to human health and longevity to life span. Ironically, usage of antibiotics to counter the various diseases caused by pathogenic microbes has been proving quite ineffective during the last decade or so. The situation is turning alarmingly grave because microbes have become increasingly resistant to almost all the antibiotics including those which had been approved quite recently [1, 2]. Pharmaceutical companies find it uneconomical to be in this evolutionary race with multiple drug resistant microbes [3-6]. In fact, microbes have evolved defence mechanisms against antibiotics by exploiting their large reservoir of genetic variability [6]. The needs are to search new drug targets [7, 8] and/or attenuate bacterial pathogenicity [9] by preventing dissemination of genes conferring resistance and those responsible for their virulent characteristics [3] and design drugs accordingly.

Among the various microbial pathogens, those notorious for causing high morbidity and mortality among living beings initiate the infection process within specialized structures – biofilms, which enable microbes to withstand adverse

environmental conditions, such as the host defence components and drugs including antibiotics used for treating the disease [10, 11]. Most pathogenic microbes develop it in a density dependent manner by a process termed as quorum sensing (QS) [12], which is regulated by the signal molecules released in their environment [13]. In Gram-negative bacteria, the most widely reported QS system is LuxR/I-type, where signals such as acylated homoserine lactones (AHLs) regulate the expression of virulence factors like elastase and pycocyanin production by *Pseudomonas aeruginosa* [14], antibiotic production by *Erwinia carotovora*, pigment and antibiotic production by *Serratia liquefaciens* [15]. The AHL-type pheromones differ in their specificity due to variations in acyl chain length, degree of saturation and substitutes at the third carbon of the lactone ring [16]. More than 70 bacterial species produce AHL type QS [15, 17].

In contrast to AHL mediated QS, which allows bacteria to compete and dominate the bacterial community, there are others which have developed mechanisms to degrade these signals. The process of interference with QS, known as quorum quenching (QQ), has been reported among prokaryotes and a few eukaryotic organisms [18, 19]. Bacteria are known to produce AHL-degrading enzymes such as AHL-lactonases and AHL-acylases. Lactonases hydrolyze the lactone ring of AHL and are found in numerous *Bacillus* spp. [20, 21]. Heterologous expression of *aiiA* (encoding for AHL-lactonase) from *Bacillus* spp. in *Pseudomonas*, *Burkholderia* and *Erwinia* resulted in marked decrease in their QS signal

*Address correspondence to this author at the Microbial Biotechnology and Genomics, Institute of Genomics and Integrative Biology (IGIB), CSIR, Delhi University Campus, Mall Road, Delhi-110007, India; Tel: +911127666156, 27666157 Ext 132; Fax: +911127667471; E-mail: vckalia@igib.res.in, vc_kalia@yahoo.co.in

sequences [43-45]. MEME searches can be performed via the web server (<http://meme.nbcr.net>) and its mirror sites [44]. The same web server also allows access to motif alignment and search tool to search sequence databases for matches to motifs. To successfully discover motifs with MEME, it is necessary to choose and prepare the input sequences carefully. Ideally, the sequences should be <1000 base pairs long [46].

Data sets in FASTA format were submitted group wise in MEME program Version 4.0.0 (<http://meme.nbcr.net/meme4/cgi-bin/meme.cgi>). In order to obtain maximum number of motifs in our sequences, we modified default settings from 3 motifs to 10 motifs. MEME search stops when this number of motifs has been found, or when none can be found with E-value less than 10000 (<http://meme.nbcr.net/meme4/meme-input.html#width>). We used default setting zero or one motif per sequence to get the occurrence of single motif which is distributed among the sequences. The default value of motif widths, set between 6 (minimum) and 50 (maximum) were modified and re-set between 25 and 30, respectively. Each of the 10 signatures (25 to 30 nucleotides long) (Table S2 to S17) was checked for its frequency of occurrence among all the sequences of a particular taxonomic group and the ones with highest frequency and did not appear in other taxonomic groups were considered unique to a particular reference taxonomic group.

In our analysis, sequences from each taxonomic group were analyzed for signature using MEME [43]. The MEME result gives a graphic representation and Motifs in regular expression format. MEME analysis were conducted for each taxa like Actinobacteria, Firmicutes, α -, β -, γ -, and δ -Proteobacteria, Ascomycota, Bacteroidetes, Euryarchaeota. The number of sequences of γ -Proteobacteria with reference to acylase from *Ralstonia* sp. Table were too large to be accommodated on line by the MEME Software (Word limit up to 60,000). In this case, we segregated the members of γ -bacterial group and made three case studies and searched for the signatures in i) *Pseudomonas* spp., ii) *Shewanella* spp. and iii) rest of the organisms (Designated as Others). The signatures obtained from the MEME program were cross-checked in other Tables and the other taxa in the same Table using online tool Sequence Manipulation Suite SMS [47].

VALIDATION OF SIGNATURES AND REGULAR EXPRESSIONS

The unique signatures - obtained after crosschecking with other taxonomic groups - were used as query sequence to BLAST against the sequenced microbial genomes available in NCBI database (<http://www.ncbi.nlm.nih.gov/>), to validate them [45]. Phylogenetic trees based on complete sequence of the gene and the strings based on the signatures were compared for AHL-lactonase among Firmicutes and AHL-acylase among γ -Proteobacteria - *Pseudomonas* spp. For developing strings, each of the 10 signatures was represented by a single alphabet - A to J.

RESULTS

Quorum sensing phenomenon has been reported from a wide range of organisms, which employ these primarily to sense environmental stress by releasing signal molecules: i)

AHL by Gram-negative bacteria and ii) oligopeptides by Gram-positive bacteria. In contrast, quite a few organisms also have the abilities to degrade these signal molecules to ensure their survival. Here we are presenting the phylogenetic diversity of AHL inactivating enzymes: AHL-lactonases and AHL-acylases, which act by hydrolyzing the lactone ring and by cleaving the acyl chain, respectively.

PHYLOGENY OF AHL-LACTONASE AND AHL-ACYLASE

Using *Bacillus* sp. SB4 AHL-lactonase (Accession No. AAR854821) (Fig. 1) amino acid sequences as reference, BLAST resulted in detecting 110 sequences in different organisms showing completely conserved domains (Table S1). Sequences for this enzyme were found to be distributed among organisms belonging to Actinobacteria, Acidobacteria, Bacteroidetes, Chloroflexi, Deinococcus-Thermus, Firmicutes, α -Proteobacteria, β -Proteobacteria, γ -Proteobacteria, δ -Proteobacteria, Euryarchaeota, Crenarchaeota, Sphingobacteria, Spirochaetales, Nitrospirales, and Planctomycetes. Among these, two major groups of organisms possessing AHL-lactonase were represented by Firmicutes - 12 genera and 22 spp. and α -Proteobacteria - 18 genera and 22 spp. (Table S1).

On the other hand, with *Ralstonia* sp. XJ12B AHL-acylase (Accession No: AAO41113.1) (Fig. 1) as reference, BLAST resulted in detecting 94 sequences of different organisms showing completely conserved domains (Table S1). The distribution of sequences for AHL-acylase was rather limited in comparison to AHL-lactonase. It was found among Actinobacteria, Cyanobacteria, Bacteroidetes, Deinococcus-Thermus, Firmicutes, α -Proteobacteria, β -Proteobacteria, γ -Proteobacteria, δ -Proteobacteria, Euryarchaeota and Crenarchaeota. The distribution pattern of this enzyme was in clear contrast to that of AHL-lactonase. The largest group of organisms (59) possessing AHL-acylase belonged to γ -Proteobacteria - 20 genera and 31 spp. Here, Firmicutes and α -Proteobacteria were among the smaller groups constituted by 3 and 10 genera, respectively. The frequency of occurrence of AHL-acylase was relatively low among other Proteobacteria. Another contrasting scenario was observed among cyanobacterial members, which were found to possess only AHL-acylase. On the other hand, organisms belonging to Acidobacteria, Sphingobacteria, Spirochaetales, Nitrospirales and Planctomycetes were found to possess only AHL-lactonase (Table S1).

Multiple sequence alignment of all the sequences revealed that conserved regions for gene encoding for AHL-lactonase and AHL-acylase are located at the positions ranging i) from 467 to 484, 557 to 571 and 737 to 755 and ii) from 236 to 254 and 1892 to 1907 nts, respectively (Table 1). These regions can be used for designing 'universal' primers for amplifying genes from diverse taxonomic classes. In addition, certain conserved regions were also observed within each taxonomic group, which can be exploited for amplifying genes specific to them. The following conserved sequences can be used for amplifying lactonase: i) Firmicutes - TA[CT]ATTA at positions 535 to 542 nts, and ATATCGT at 774 to 780 nts, ii) β -Proteobacteria - CGCA-GACCA at 681 to 689 nts, and CGG[AC]CG at 521 to 523

nts, etc. Similar regions were also observed among γ -Proteobacteria, δ -Proteobacteria and Ascomycota for lactonase gene (Table 1) and acylase type gene among Actinobacteria, Cyanobacteria, α -Proteobacteria, β -Proteobacteria and Bacteroidetes (Table 1).

PHYLOGENETIC DISCREPANCIES

The phylogenetic trees (Figs. 2-5) based on organisms found to possess AHL-lactonase and AHL-acylase followed the taxonomic distribution, which is expected on the basis of

Table 1. Location of Conserved Sequences in AHL-Lactonase and AHL-Acylase Genes of Taxonomically Diverse Bacterial Groups

| Taxonomic Group | Position (nts) ^a | Conserved Regions |
|---------------------------|-----------------------------|-------------------------------------|
| AHL-lactonase gene | | |
| All ^b | 467-484 | TGGTC[AG]AAATGCAGATGGGT |
| | 557-571 | ATTCGACCATGCCG |
| | 737-755 | AGCGATCACGCGTGTCC |
| Actinobacteria | 560-570 | TCGACCACG[CG]CG |
| Firmicutes | 535-542 | TA[CT]ATTA |
| | 559-566 | TTCGATCA |
| | 774-780 | ATATCGT |
| Alphaproteobacteria | 562-571 | GACCAT[ACG]TCGT |
| Betaproteobacteria | 521-523 | CGG[AC]CG |
| | 681-689 | CGCAGACCA |
| Gammaproteobacteria | 527-534 | ATGTCGAG |
| | 548-552 | CCCAT |
| Deltaproteobacteria | 518-522 | A[GT]CCC |
| Euryarchaeota | 560-569 | T[CT]GACCACGC |
| Ascomycota | 338-347 | TTGAXCATCC |
| | 520-539 | ATCAAAGACGTCAAGGCAGT |
| | 550-571 | CA[CT]TTACATCTCGA[CT]CACGCAG |
| | 628-633 | CACGAG |
| AHL-acylase gene | | |
| All ^c | 236-254 | TGATCGCCGAAATCCGGCG |
| | 1892-1907 | GTATGGGTCCAGGCCA |
| Actinobacteria | 704-713 | TACCGGGTCG |
| | 1177-1181 | AC[AC]C[CG]GTCA[CG]C |
| | 1208-1217 | CCA[CG]GAGCTG |
| Cyanobacteria | 1522-1527 | TTGAAA |
| Alphaproteobacteria | 486-495 | TCGATGCG |
| Betaproteobacteria | 328-340 | GCGCAGGCAACC |
| Gammaproteobacteria | 309-320 | CGGCATCGGTTA |
| | 586-595 | GGCTTCAACC |
| Bacteroidetes | 1490-1497 | TC[CG]GCATA |

^a Nucleotides in the gene from 5' to 3'.

^b Gene sequences from taxonomic groups listed under AHL-lactonase gene.

^c Gene sequences from taxonomic groups listed under AHL-acylase gene.

Fig. (2). Phylogenetic tree (Upper half) based on protein sequences of organisms containing conserved domains with respect to *Bacillus* sp. SB4 AHL lactonase as reference (For Taxonomic details refer to Additional file 3). A Neighbor-Joining analysis with Jukes-Cantor correction and bootstrap support was performed on the protein sequences. Bootstrap values are given at nodes, 1000 bootstrap replicates were run. The leaf contains Accession number and name of the organism.

Fig. (3). Phylogenetic tree (Lower half) based on protein sequences of organisms containing conserved domains with respect to *Bacillus* sp. SB4 AHL lactonase as reference (For Taxonomic details refer to Additional file 3). A Neighbor-Joining analysis with Jukes-Cantor correction and bootstrap support was performed on the protein sequences. Bootstrap values are given at nodes, 1000 bootstrap replicates were run. The leaf contains Accession number and name of the organism.




Fig. (4). Phylogenetic tree (Upper half) based on protein sequences of organisms containing conserved domains with respect to *Ralstonia* sp. XJ12B AHL acylase as reference (For Taxonomic details refer to Additional file 3). A Neighbor-Joining analysis with Jukes-Cantor correction and bootstrap support was performed on the protein sequences. Bootstrap values are given at nodes, 1000 bootstrap replicates were run. The leaf contains Accession number and name of the organism.




Fig. (5). Phylogenetic tree (Lower half) based on protein sequences of organisms containing conserved domains with respect to *Ralstonia* sp. XJ12B AHL acylase as reference (For Taxonomic details refer to Additional file 3). A Neighbor-Joining analysis with Jukes-Cantor correction and bootstrap support was performed on the protein sequences. Bootstrap values are given at nodes, 1000 bootstrap replicates were run. The leaf contains Accession number and name of the organism.

their 16S rDNA sequences. A few phylogenetic discrepancies were observed for AHL-lactonase (Figs. 2,3, Table 2), the notable being the associations between i) *Moorella thermoacetica* ATCC 39073 (Firmicutes) and *Burkholderia graminis* C4D1M (β -Proteobacteria) (Boot Strap Value, BV 1000), ii) Actinobacteria (*Mycobacterium* sp. MCS and *Rubrobacter xylanophilus* DSM9941) and members of α -Proteobacteria (*Granulibacter bethesdensis* CGDNIH1, *Acidiphilum cryptum* Jf-5 and *Agrobacterium tumefaciens*) (BV 820), and iii) *Deinococcus radiodurans* R1 (Deinococcus-Thermus) and *Xylella fastidiosa* (γ -Proteobacteria) (BV 1000). Similarly, certain taxonomic discrepancies were also recorded in the phylogenetic tree (Figs. 4,5, Table 2) drawn with respect to sequences of organisms possessing AHL-acylase: i) *Solibacter usitatus* Ellin 6076 (Acidobacteria) and *Plesiocystis pacifica* SIR-1 (δ -Proteobacteria) (BV 1000), ii) *D. radiodurans* R1 (Deinococcus-Thermus) and *Ralstonia metallidurans* CH34, *R. solanacearum* GMI1000 and *Ralstonia* sp. XJ12B (β -Proteobacteria (BV 1000). Since these associations between widely distributed organisms show very high BVs, these may be interesting cases of horizontal gene transfers.

ORGANISMS POSSESSING GENES FOR AHL-LACTONASE AND AHL-ACYLASE

Sixteen bacterial strains of 5 different genera and a group of 3 strains belonging to marine gamma proteobacterium

were found to possess genes for both the enzyme types: AHL-lactonase and AHL-acylase (Table 3). The diversity of these 19 strains was evident by their distribution among 5 different taxonomic groups: i) *Streptomyces* spp. (Actinobacteria), ii) *Deinococcus* spp. (Deinococcus-Thermus), iii) *Hyphomonas* sp. (α -Proteobacteria), iv) *Ralstonia* spp. (β -Proteobacteria), v) *Photorhabdus* sp. (γ -Proteobacteria), vi) marine gamma proteobacterium. Three out of these 19 strains can be more clearly classified as those which possessed genes for both the enzyme types: i) *D. radiodurans* R1, ii) *Hyphomonas neptunium* ATCC 15444 and iii) *Photorhabdus luminescens* subsp. *laumondii* TTO1. This observation is supported by the presence motifs for lactonase and acylase in these strains.

SIGNATURE ANALYSIS – FREQUENCY AND DISTRIBUTION PATTERN

Ten signatures (nucleotides) were obtained through MEME for different taxonomic groups with reference to: i) AHL-lactonase gene (*aiiA*) of *Bacillus* sp. SB4 Accession No. AAR85482.1 (Figs. S1 to S8 and Tables S2 to S9) and ii) AHL-acylase gene (*aiiD*) of *Ralstonia* sp. XJ12B Accession No. AAO41113.1 (Figs. S9 to S19 and Tables S10 to S17). (See Additional File 4 for a description of data in Tables S2 to S17 and Figs. S1 to S17). Nucleotide signatures found to be present at high frequency within the taxonomic group and absent from all other groups were initially classified as

Table 2. Potential Cases of Horizontal Gene Transfer of Acyl-Homoserine Lactone-Acylase and –Lactonase

| Incongruent Phylogenetic Associations | | Boot Strap Value |
|---|--------------------------|------------------|
| Organism(s) | Taxonomy | |
| <i>Bacillus</i> sp. SB4 (AHL-lactonase)^a | | |
| <i>Moorella thermoacetica</i> ATCC 39073 | Firmicutes | 1000 |
| <i>Burkholderia graminis</i> C4D1M | β -Proteobacteria | |
| <i>Mycobacterium</i> sp. MCS | Actinobacteria | 820 |
| <i>Rubrobacter xylanophilus</i> DSM9941 | | |
| <i>Granulibacter bethesdensis</i> CGDNIH1 | | |
| <i>Acidiphilum cryptum</i> Jf-5 | α -Proteobacteria | |
| <i>Agrobacterium tumefaciens</i> | | |
| <i>Deinococcus radiodurans</i> R1 | Deinococcus-Thermus | |
| <i>Xylella fastidiosa</i> | γ -Proteobacteria | |
| <i>Ralstonia</i> sp. XJ12B (AHL-acylase)^a | | |
| <i>Solibacter usitatus</i> Ellin 6076 | Acidobacteria | 1000 |
| <i>Plesiocystis pacifica</i> SIR-1 | δ -Proteobacteria | |
| <i>Deinococcus radiodurans</i> R1 | Deinococcus-Thermus | 1000 |
| <i>Ralstonia metallidurans</i> CH34 | β -Proteobacteria | |
| <i>R. solanacearum</i> GM1000 | | |
| <i>Ralstonia</i> sp. XJ12B | | |

^a Reference organism.

Table 3. Taxonomic Distribution of Organisms (Genera) Possessing Enzymes Acyl-Homoserine Lactone-Acylase as well as – Lactonase

| AHL-Lactonase | | AHL-Acylase | |
|--|--------------------|---|---|
| <i>Bacillus</i> sp. SB4 ^a (GenBank AAR85482.1) | Motif ^b | <i>Ralstonia</i> sp. XJ12B ^a (GenBank AAO41113.1) | Motif ^c |
| Actinobacteria | | | |
| <i>Streptomyces clavuligerus</i> ATCC 27064 | + | <i>Streptomyces</i> sp. FERM BP-5809 | + (I ₅₄ -D ₆₁) |
| | | <i>Streptomyces</i> sp. M664 | + (I ₅₄ -D ₆₁) |
| | | <i>Streptomyces</i> sp. Mg1 | + (I ₅₄ -D ₆₁) |
| | | <i>Streptomyces griseus</i> subsp. <i>griseus</i> NBRC 13350 | + (I ₅₄ -D ₆₁) |
| | | <i>S. sviveus</i> ATCC 29083 | + (L ₅₄ -D ₆₁) |
| Deinococcus-Thermus | | | |
| <i>Deinococcus radiodurans</i> R1 | + | <i>Deinococcus radiodurans</i> R1 | + (I ₅₈ -S ₆₅) |
| <i>D. geothermalis</i> DSM 11300 | + | | |
| Alphaproteobacteria | | | |
| <i>Hyphomonas neptunium</i> ATCC 15444 | + | <i>Hyphomonas neptunium</i> ATCC 15444 | + (I ₅₅ -S ₆₂) |
| Betaproteobacteria | | | |
| <i>Ralstonia eutropha</i> JMP134 | + | <i>Ralstonia</i> sp. XJ12B | + (I ₄₉ -D ₅₆), + (I ₅₂ -S ₅₉) |
| | | <i>Ralstonia eutropha</i> H16 | + (I ₆₂ -D ₆₉) |
| | | <i>R. metallidurans</i> CH34 | + (I ₅₁ -S ₅₈) |
| | | <i>R. pickettii</i> 12J | + (I ₅₁ -D ₅₈) |
| | | <i>R. solanacearum</i> GMI1000 | + (I ₅₀ -D ₅₇) |
| Gammaproteobacteria | | | |
| <i>Photorhabdus luminescens</i> subsp. <i>laumondii</i> TTO1 | + | <i>Photorhabdus luminescens</i> subsp. <i>laumondii</i> TTO1 | + (I ₅₇ -S ₆₄) |
| marine gamma proteobacterium HTCC2080 | + | marine Gamma proteobacterium HTCC2143 | + (I ₄₃ -D ₅₃) |
| | | marine Gamma proteobacterium HTCC2207 | + (L ₁₀₆ -D ₁₁₃) |

^a Reference organisms^b Motif for AHL-lactonase: His¹⁰⁶-X-Asp¹⁰⁸-His¹⁰⁹-59X-His¹⁶⁹-21X-Asp¹⁹¹^c Motif for AHL-acylase: amino acid – Ile (I) /Leu (L)⁵⁰ and Ser(S) /Asp (D)⁵⁷

unique. The BLAST (against the NCBI database) result validated the uniqueness of these signatures (Table 4). A high correlation was recorded for signatures M1 and M10 of *Bacillus* sp. SB4 and their occurrence among a large number of organisms possessing AHL-lactonase (BLAST hits with high frequency). Similarly, among the different signatures identified within each taxonomic group for AHL-acylase, those retrieved from *Pseudomonas* spp. within γ -Proteobacteria gave the BLAST hits with high frequency for organisms of this taxonomic group. Sequence homology searches using BLAST for all other signatures showed best hits for the query sequence from the respective taxonomic group but their frequency among the top 50 hits was quite low.

DISCUSSION

Bacteria with Ability to Quench QS Signals

With the information available in the literature on the presence of AHL-lactonase and AHL-acylase activities in *R.*

erythropolis W2 [30-32], we also looked for certain other organisms with potential to produce both these enzyme types. Our *in silico* study allowed us to identify organisms from five different genera with potential to produce both types of AHL-degrading enzymes. The three strains i) *D. radiodurans* R1, ii) *H. neptunium* ATCC 15444 and iii) *P. luminescens* subsp. *laumondii* TTO1 are among the top contenders for producing both the enzyme type from an individual strain. However, more such strains can be tracked among strains belonging to *Streptomyces* spp. (Actinobacteria), ii) *Ralstonia* spp. (β -Proteobacteria), and marine gamma proteobacterium. In order to reduce the efforts needed to search organisms, conserved regions can be used for designing primers for amplifying these genes. Signature sequences deduced in this study can be used as an additional tool to assign these sequences to respective taxonomic groups, at least in some cases.

Another interesting features of this *in silico* study is that *R. solanacearum* GMI1000 (β -Proteobacteria) has been

Table 4. Characteristics of Unique Nucleotide Signatures for Acyl-Homoserine Lactone-Acylase and -Lactonase Showing Closest Match to the Respective Taxonomic Group

| Taxonomic Group | Signature | | Source Organism |
|--|---|-----|---|
| | Nucleotide Sequence | No. | |
| <i>AHL-lactonase with reference to Bacillus sp. SB4</i> | | | |
| Actinobacteria | CGTGAGCTCGAGTACGGCCTGTCAATCAC ^a | M8 | <i>Mycobacterium</i> sp.MCS [gi 108796981:1991857-1992693] |
| Firmicutes | TCACTTACATTTTGATCATGCAGGAGGAAA | M1 | <i>Bacillus</i> sp. SB4 [gi 40388446 gb AY483161.1] |
| | TCCGGTTCAGTTTTATTAACGATTGATGCA | M10 | |
| Alphaproteobacteria | GTTGTGCAGTCGCACCTGCATCTCGATCAT ^a | M1 | <i>Agrobacterium tumefaciens</i> [gi 17223781:3565-4356] |
| | ATCCGCAAGGACTTCGACAAGCCGGGCT ^a | M4 | |
| Betaproteobacteria | GTGGATACGGAAGACGGCGTGGTGTGAT ^a | M7 | <i>Burkholderia graminis</i> [gi 170694799:130043-130822] |
| Gammaproteobacteria | AATCCGGTATCAATCAGTACAAGCCCTCGA ^a | M2 | <i>Acinetobacter baumannii</i> AYE [gi 169794206:627842-628654] |
| Deltaproteobacteria | ATGGTCGTTATGCAGGTGCGTGTGTATGAC ^a | M1 | <i>Desulfovibrio desulfuricans</i> G20 [gi 78355047:1350539-1351285] |
| Ascomycota | TACACGAGGAAGAGTTCAAGCATGCATGCT ^a | M2 | <i>Aspergillus oryzae</i> RIB40 [gi 169784499 ref XM_001826659.1] |
| | GATCAATTCCATGTGCGGGAGAACTACGAG ^a | M3 | |
| Euryarchaeota | <i>All signatures^a</i> | | |
| <i>AHL-acylase with reference to Ralstonia sp. XJ12B</i> | | | |
| Actinobacteria | CGGGTTCGCCCGGGGTACAACGCGTGGAT ^a | M2 | <i>Streptomyces</i> sp. M664 [gi 49781326:185-2599] and <i>Streptomyces</i> sp. FERM BP-5809 [gi 40786372:948-3362] |
| | GTCATCCGGTACACGGAGTACGGCATTCCG ^a | M3 | |
| Bacteroidetes | ATCGAGATGCTTGAAGGGGATTCCTCGA ^a | M3 | <i>Salinibacter ruber</i> DSM 13855 [gi 83814055:1438503-1440524-] |
| | GGCTTCGGGGGGAGATGGTCTTTTTG ^a | M7 | |
| | TTTGACGAGATGGAGGGCGGG ^a | M8 | |
| Cyanobacteria | GCTGGTTCCAATGCTTGGGCGATCGCACCT ^a | M5 | <i>Cyanothece</i> sp. [gi 126661348:2396-4477-] |
| Alphaproteobacteria | ATGACTCCGGCAAGCCCGAAGAAGAAAGGC ^a | M7 | <i>Erythrobacter litoralis</i> HTCC2594 [gi 85372828:319908-322139] |
| | GATATTGCCCTCGGCATCGCCATAGATGAA ^a | M10 | |
| Betaproteobacteria | CCAGCCGATGCGCAGGGTTTCTGGGTCC ^a | M3 | <i>Ralstonia eutropha</i> H16 [gi 113866031:2083444-2085906] |
| | TCCGCACGCGCAGGTTGAACAGCGGCGACA ^a | M7 | |
| Gammaproteobacteria: <i>Pseudomonas</i> spp. | CTGCCGGTGGTCAACATCGGCTTCAGCCGC | M1 | <i>Pseudomonas aeruginosa</i> PA7 [gi 152983466:2955521-2957803] |
| | TTCGTACAGAACTCCAACGACAGCGCCTGG | M2 | |
| | GATGCGTTTCTACCAGATGCACCTGACCAT | M3 | |
| | ATGCTGCTGGCCAACCCGCATTTCCCCTGG | M4 | |
| | CGGCATCGGCTACGCCTACGCGCGGACA | M6 | |
| | GCGCGATGCCAACCTGGAGAACACCCGGGT | M7 | |
| | CCGCTGGACCGCTATGGCGTGCCGCACAT | M9 | |
| | ATCCCCTGGGTCAACACCCTGGCCGCGGA | M10 | |

Table 4. contd....

| Taxonomic Group | Signature | | Source Organism |
|--|---|-----|--|
| | Nucleotide Sequence | No. | |
| <i>AHL-acylase with reference to Ralstonia sp. XJ12B</i> | | | |
| Gammaproteobacteria: <i>Shewanella</i> spp. | CAGATAATAAAGGCAACGCCTTCTACATCG ^a | M1 | <i>Shewanella woodyi</i> ATCC 51908 [gij170724370:166858-169239] |
| | CCCTTTGAGCGTGCCCCAAAGTGATGCGT ^a | M2 | |
| | GGCCGTGATAAAACCGAATCTGGTAGCGGT ^a | M3 | |
| | TTAATCGGCTTCAATGAACATGTTGCTTGG ^a | M6 | |
| | GGTTATGGGCAAGCCTATGCCCATGCTCAG ^a | M10 | |
| Others Gammaproteobacterial members | TTATCCGCTTCGATGTGCGGCGCGCCGAA ^a | M8 | <i>Hahella chejuensis</i> KCTC 2396 [gij83642913:5633846-5636326] |

^a Sequence homology searches using BLAST for these signatures showed best hits for the query sequence but the frequency among the top 50 hits was quite low.

found to possess gene and motif for AHL-acylase. This observation finds support from the work which demonstrates the presence of AHL-acylase in *R. solanacearum* GMI1000, through cloning and expression of *aac* gene [25]. It led to inactivation of four different AHLs [25]. Another interesting observation was the presence of genes for quenching the QS signals in archaeal member *A. fulgidus*. Although archaea have not been reported so far to be directly involved in causing infectious diseases [48], however, there are evidences of their presence in endodonic infections [49]. The presence of these genes support the possibility of their role in infections.

MOTIFS OF AHL-LACTONASES AND AHL-ACYLASES

AHL-lactonase is a member of metallo-hydrolase superfamily containing the motif: His¹⁰⁴-X-His¹⁰⁶-X-Asp¹⁰⁸-His¹⁰⁹, which resembles zinc binding motif of several other metallo-enzymes such as glyoxalase II, aryl sulfatase and β -lactamase. The motif His¹⁰⁶-X-Asp¹⁰⁸-His¹⁰⁹-59X-His¹⁶⁹-21X-Asp¹⁹¹ has been reported to be essential for AHL-lactonase activity [18, 22].

A very interesting scenario was observed while searching for motifs essential for AHL-lactonase enzyme activities. It was observed to be that motif essential for AHL-lactonases - His¹⁰⁶-X-Asp¹⁰⁸-His¹⁰⁹-59X-His¹⁶⁹-21X-Asp¹⁹¹ was present in all the organisms having potential for AHL-lactonase except *R. solanacearum* GMI1000, *A. baumannii* AB0057, *A. baumannii* ACICU, *A. baumannii* AYE, *Acinetobacter* sp. ADP1 and *Pseudomonas stutzeri* A1501 (Table 3). Incidentally, these latter 6 bacteria lacked even the motif specific for metallo-enzymes in general- His¹⁰⁴-X-His¹⁰⁶-X-Asp¹⁰⁸-His¹⁰⁹. Among the organisms found to possess both the motifs – all belonged to *Bacillus* spp.: *B. anthracis* str. Ames, *B. cereus*, *B. cereus* AH1134, *B. cereus* AH820, *B. cereus* ATCC 10987, *B. cereus* ATCC 14579, *B. cereus* B4264, *B. cereus* E33L, *B. cereus* G9241, *B. cereus* NVH0597-99, *Bacillus* sp. 240B1, *Bacillus* sp. 42, *Bacillus* sp. 91, *Bacillus* sp. A24, *Bacillus* sp. COT1, *Bacillus* sp. CSX-1, *Bacillus* sp. SB4, *B.*

subtilis, *B. thuringiensis*, *B. thuringiensis* serovar *alesti*, *B. thuringiensis* serovar *canadensis*, *B. thuringiensis* serovar *galleriae*, *B. thuringiensis* serovar *israelensis* ATCC 35646, *B. thuringiensis* serovar *jinhongiensis*, *B. thuringiensis* serovar *kim*, *B. thuringiensis* serovar *konkukian* str. 97-27, *B. thuringiensis* serovar *kurstaki*, *B. thuringiensis* serovar *kyu-shuensis*, *B. thuringiensis* serovar *ostrinae*, *B. thuringiensis* serovar *oswaldocruzi*, *B. thuringiensis* serovar *pakistanii*, *B. thuringiensis* serovar *toumanoffi* and *B. weihenstephanensis* KBAB4.

Available literature reveals that AHL-acylases are structurally similar with respect to the two amino-acids at positions 50 and 57 [18] as a) AiiD of *Ralstonia* sp. XJ12B has Ile⁵⁰ and Ser⁵⁷ [26], b) PvdQ of *P. aeruginosa* PAO1 has Leu⁵⁰ and Asp⁵⁷ [27], and c) AhlM of *Streptomyces* sp. has Leu⁵⁰ and Ser⁵⁷ [50]. These share features with Ntn hydrolases including the signal peptide, α - and β -subunits [51].

On the other hand, Cephalosporin-acylase has Gln and Arg at positions 50 and 57 respectively [18]. Our analysis reveals that most of the organisms had amino-acids of AHL acylase in a slightly different positions than those reported in the literature (Table 3). Unique features were observed in i) *Streptomyces* spp., where the amino acids of motif for acylase resembled either those found in PvdQ or were a combination of Ile of AiiD and Asp of PvdQ, ii) *Ralstonia* spp., where Ser was substituted with Asp and also that *Ralstonia* sp. XJ12B was found to possess two motifs – a) Ile⁵² and Ser⁵⁹ and b) Ile⁴⁹ and Asp⁵⁶. In view of the fact that in spite of such a large number of organisms, which had the conserved domains for AHL-acylase, only few seem to have retained the property observed at respective positions as reported in literature [18, 26, 27, 50]. It may not be too inappropriate to conclude that amino-acids at positions 50 and 57 might have changed during the course of evolution.

A few other organisms which deserve attention are: i) *Burkholderia* spp. (β -Proteobacteria), which had motifs for

lactonase and cephalosporin acylase, ii) *Pseudomonas* sp. (γ -Proteobacteria), which possessed two motifs for AHL-acylase and one for cephalosporin acylase, and iii) *A. baumannii* AYE (γ -Proteobacteria), which showed motifs for cephalosporin acylase.

USING SIGNATURES TO DESIGN PRIMERS

Bacillus spp. and *Pseudomonas* spp. have been reported to possess repeating elements in the 16S rDNA [2, 45]. We observed such repeats in the AHL-lactonase and -acylase gene in different members of these two genera. Individually, these signatures were validated by using them as query sequences in BLAST search. Since the frequency of repeating elements in this gene was quite high in the case of *Pseudomonas* spp., we also checked them for regular expressions [52]. We observed a very good co-relation between the full length gene sequence and the strings (Fig. S18a, b). The nucleotide signatures identified in our study for each taxonomic group may enable one to design primers to amplify genes for AHL-lactonase [20, 24, 53] or AHL-acylase and identify a possible taxonomic group within a microbial community.

THE ROLE OF QUORUM QUENCHERS

The role of QQ enzymes *in vivo* is debatable [54]. They may be involved in metabolic pathways where AHL and its products may be used as energy and carbon sources -such as by *Variovorax paradoxus* and *Arthrobacter* spp. [55-57]. AHL-acylase PvdQ from *P. aeruginosa* is thought to be involved in biosynthesis of protein (pyoverdine) necessary for acquisition of iron [28, 54, 58]. The role of the enzymes - lactonase and acylase for QQ has been proposed to prevent food, prevent biofilm growth, surface fouling, treatment of bacterial infections, etc. [59]. It can act as a potential drug, which will selectively block the virulence and pathogenic traits [60]. Heterologous expression of *Bacillus* AiiA lactonase in *Burkholderia thailandensis* [24], in *P. aeruginosa* PAO1 [23], in *Vibrio harveyi* [53], in *E. carotovora* [22] have been found to disrupt QS mediated properties. In fact, certain antibiotics become effective on biofilms which have been exposed to AHL degrading enzymes [61], which imply a combination of the two treatments can be complementary and effective in controlling the pathogenic organism(s). Although there may be barriers to their diffusion within the microbial communities in the soil environment, however, their usage as a drug may not face similar barriers and may prove effective.

CONCLUSION

Using available metabolic and genomic databases data-bases, we have been able to trace potential organisms such as i) *Deinococcus radiodurans* R1, ii) *Hyphomonas neptunium* ATCC 15444 and iii) *Photorhabdus luminescens* subsp. *laumondii* TTO1 to possess genes for AHL-lactonases and -acylases. In addition, all the cyanobacterial members were found to possess genes only for AHL-acylase whereas members of Acidobacteria, Sphingobacteria, Spirochaetales, Nitrospirales and Planctomycetes were found to possess genes only for AHL-lactonase. It also revealed those bacteria

which evolved to produce either of the two enzymes, an indication of the environmental stresses encountered during their evolution. The presence of genes for quenching the QS signals in archaea has further increased the range of organisms which can be exploited for producing these enzymes for preventing food spoilage, bioremediation and design drugs against bacterial infections.

Phylogenetic analyses and multiple sequence alignment of the gene sequences coding for the two enzymes revealed consensus sequences, which can be used to design primers for amplifying these genes even among mixed cultures and metagenomes. Nucleotide signature sequences deduced in this study can be used as an additional tool to assign these sequences to respective taxonomic groups. A few phylogenetic discrepancies observed for AHL-lactonase and -acylase may imply HGT and help us identify those organisms which can be subjected to genetic engineering with a relative ease.

It has been proposed that since lactonase activity is not influenced by the acyl chain length of the AHL signal molecule, hydrolysis of the ring may be more effective approach for QQ. As a wide range of *Bacillus* spp. have an ability to produce lactonase, it may be the organism of choice in future. The usage of *Bacillus* is quite beneficial since it has been labeled as industrial workhorse and accorded the designation GRAS (Generally Regarded As Safe) by FDA. These organisms can be exploited for producing drugs against pathogens.

SUPPLEMENTARY MATERIAL

Supplementary material is available on the publishers web site along with the published article.

ACKNOWLEDGEMENTS

We are thankful to Directors of Institute of Genomics and Integrative Biology, CSIR; National Environmental Engineering Research Institute, CSIR grant SIP 16 for providing necessary funds, facilities and moral support.

AUTHOR'S CONTRIBUTIONS

VCK has contributed towards the conceived and designed the experiments, analyzed the data and wrote the paper.

CRS and HJP have performed the experiments, and contributed reagents/ materials/ analysis tools. All authors read and approved the final manuscript.

REFERENCES

- [1] D'Costa VM, McGrann KM, Hughes DW, Wright GD. Sampling the antibiotic resistome. *Science* 2006; 311: 374-377.
- [2] Purohit HJ, Raje DV, Kapley A Identification of signature and primers specific to genus *Pseudomonas* using mismatched patterns of 16S rDNA sequences. *BMC Bioinformatics* 2003; 4: 19.
- [3] Courvalin P. Predictable and unpredictable evolution of antibiotic resistance. *J Intern Med* 2008; 264: 4-16.
- [4] Kalia VC, Lal S, Ghai R, Mandal M, Chauhan A. Mining genomic databases to identify novel hydrogen producers. *Trends Biotechnol* 2003; 21: 152-156.
- [5] Spellberg B, Powers JH, Brass EP, Miller LG, Edwards Jr JE. Trends in antimicrobial drug development: implications for the future. *Clin Infect Dis* 2004; 38: 1279-1286.

- [6] Wright GD The antibiotic resistome: the nexus of chemical and genetic diversity. *Nat Rev Microbiol* 2007; 5: 175-186.
- [7] Cars O, Högberg LD, Murray M, *et al.* Meeting the challenge of antibiotic resistance. *Br Med J* 2008; 337: 726-728.
- [8] Mdluli K, Spigelman M. Novel targets for tuberculosis drug discovery. *Curr Opin Pharmacol* 2006; 6: 459-467.
- [9] Bjarnsholt T, Givskov M. Quorum-sensing blockade as a strategy for enhancing host defences against bacterial pathogens. *Phil Trans R Soc B* 2007; 362: 1213-1222.
- [10] Anderson GG, Palermo JJ, Schilling JD, Roth R, Heuser J, Hultgren SJ. Intracellular bacterial biofilm-like pods in urinary tract infections. *Science* 2003; 301: 105-107.
- [11] Furukawa S, Kuchma SL, O'toole GA. Keeping their options open: acute versus persistent infections. *J Bacteriol* 2006; 188: 1211-1217.
- [12] Abraham WR. Controlling biofilms of gram-positive pathogenic bacteria. *Curr Med Chem* 2006; 13: 1509-1524.
- [13] O'Toole GA, Kolter R. Initiation of biofilm formation in *Pseudomonas fluorescens* WCS365 proceeds via multiple, convergent signalling pathways: a genetic analysis. *Mol Microbiol* 1998; 28: 449-461.
- [14] Muh U, Schuster M, Heim R, Singh A, Olson ER, Greenberg EP. Novel *Pseudomonas aeruginosa* quorum-sensing inhibitors identified in an ultra-high-throughput screen. *Antimicrob Agents Chemother* 2006; 50: 3674-3679.
- [15] Henke JM, Bassler BL. Bacterial social engagements. *Trends Cell Biol* 2004; 14:648-656.
- [16] González JE, Keshavan ND. Messing with bacterial quorum sensing. *Microbiol Mol Biol Rev* 2006; 70: 859-875.
- [17] Williams P, Winzer K, Chan W, Cámara M. Look who's talking: communication and quorum sensing in the bacterial world. *Phil Trans R Soc B* 2007; 362: 1119-1134.
- [18] Dong Y-H, Wang L-H, Zhang L-H. Quorum-quenching microbial infections: mechanisms and implications. *Phil Trans R Soc B* 2007; 362: 1201-1211.
- [19] Turovskiy Y, Kashtanov D, Paskhover B, Chikindas ML. Quorum sensing: Fact, fiction, and everything in between. *Adv Appl Microbiol* 2007; 62: 191-234.
- [20] Dong Y-H, Gusti AR, Zhang Q, Xu J-L, Zhang L-H. Identification of quorum-quenching N-acyl homoserine lactonases from *Bacillus* species. *Appl Environ Microbiol* 2002; 68: 1754-1759.
- [21] Zhang L-H. Quorum quenching and proactive host defense. *Trends Plant Sci* 2003; 8: 238-244.
- [22] Dong Y-H, Xu J-L, Li XZ, Zhang L-H. AiiA, an enzyme that inactivates the acylhomoserine lactone quorum-sensing signal and attenuates the virulence of *Erwinia carotovora*. *Proc Natl Acad Sci USA* 2000; 97: 3526-3531.
- [23] Reimann C, Ginet N, Michel L, *et al.* Genetically programmed autoinducer destruction reduces virulence gene expression and swarming motility in *Pseudomonas aeruginosa* PAO1. *Microbiology* 2002; 148: 923-932.
- [24] Ulrich RL. Quorum quenching: Enzymatic disruption of N-Acylhomoserine lactone-mediated bacterial communication in *Burkholderia thailandensis*. *Appl Environ Microbiol* 2004; 70: 6173-6180.
- [25] Chen C-N, Chen C-J, Liao C-T, Lee C-Y. A probable aculeacin A acylase from the *Ralstonia solanacearum* GMI1000 is N-acyl-homoserine lactone acylase with quorum-quenching activity. *BMC Microbiol* 2009; 9: 89.
- [26] Lin YH, Xu JL, Hu J, Wang LH, Ong SL, Leadbetter JR, Zhang LH. Acyl-homoserine lactone acylase from *Ralstonia* strain XJ12B represents a novel and potent class of quorum-quenching enzymes. *Mol Microbiol* 2003; 47: 849-860.
- [27] Huang JJ, Han J-I, Zhang L-H, Leadbetter JR. Utilization of acyl-homoserine lactone quorum signals for growth by a soil pseudomonad and *Pseudomonas aeruginosa* PAO1. *Appl Environ Microbiol* 2003; 69: 5941-5949.
- [28] Lamont IL, Martin LW. Identification and characterization of novel pyoverdine synthesis genes in *Pseudomonas aeruginosa*. *Microbiology* 2003; 149: 833-842.
- [29] Sio CF, Otten LG, Cool RH, *et al.* Quorum quenching by an N-Acyl-Homoserine lactone acylase from *Pseudomonas aeruginosa* PAO1. *Infect Immun* 2006; 74: 1673-1682.
- [30] Uroz S, D'Angelo-Picard C, Carlier A, *et al.* Novel bacteria degrading N-acylhomoserine lactones and their use as quenchers of quorum-sensing-regulated functions of plant-pathogenic bacteria. *Microbiology* 2003; 149: 1981-1989.
- [31] Uroz S, Oger PM, Chapelle E, Adeline MT, Faure D, Dessaux Y. A *Rhodococcus qsdA*-encoded enzyme defines a novel class of large-spectrum quorum-quenching lactonases. *Appl Environ Microbiol* 2008; 74: 1357-1366.
- [32] Uroz S, Chhabra SR, Cámara M, Williams P, Oger PM, Dessaux Y. N-Acylhomoserine lactone quorum-sensing molecules are modified and degraded by *Rhodococcus erythropolis* W2 by both amidolytic and novel oxidoreductase activities. *Microbiology* 2005; 151: 3313-3322.
- [33] Kalia VC, Chauhan A, Bhattacharyya G, Rashmi. Genomic databases yield novel bioplastic producers. *Nat Biotechnol* 2003; 21: 845-846.
- [34] Kalia VC, Rani A, Lal S, Cheema S, Raut CP. Combing databases reveals potential antibiotic producers. *Expert Opin Drug Discov* 2007; 2: 211-224.
- [35] Purohit HJ, Cheema S, Lal S, Raut CP, Kalia VC. In search of drug targets for *Mycobacterium tuberculosis*. *Infect Dis Drug Targets* 2007; 7: 245-250.
- [36] Marchler-Bauer A, Bryant SH. "CD-Search: protein domain annotations on the fly". *Nucleic Acids Res* 2004; 32(W): 327-331.
- [37] Marchler-Bauer A, Anderson JB, Chitsaz F, *et al.* "CDD: specific functional annotation with the Conserved Domain Database". *Nucleic Acids Res* 2009; 37(D): 205-210.
- [38] Saitou N, Nei M. The Neighbor-joining Method: A new method for reconstructing phylogenetic trees. *Mol Biol Evol* 1987; 4: 406-425.
- [39] Larkin MA, Blackshields G, Brown NP, *et al.* Clustal W and Clustal X version 2.0. *Bioinformatics* 2007; 23: 2947-2948.
- [40] Roderic DMP. TreeView: An application to display phylogenetic trees on personal computers. *Bioinformatics* 1996; 12: 357-358.
- [41] Clamp M, Cuff J, Searle SM, Barton GJ. The Jalview Java Alignment Editor. *Bioinformatics* 2004; 20: 426-427.
- [42] Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ. Jalview Version 2 - a multiple sequence alignment editor and analysis workbench. *Bioinformatics* 2009; 25: 1189-1191.
- [43] Bailey TL, Elkan C. Fitting a mixture model by expectation maximization to discover motifs in biopolymers. In: Altman RB, Brutlag DL, Karp PD, Lathrop RH, Searls DB, eds. *Proceedings of the Second International Conference on Intelligent Systems for Molecular Biology*. AAAI Press, Menlo Park, CA, 1994; pp. 28-36.
- [44] Bailey TL, Williams N, Misleh C, Li WW. MEME: discovering and analyzing DNA and protein sequence motifs. *Nucleic Acids Res* 2006; Web Server issue: W369-W373.
- [45] Porwal S, Lal S, Cheema S, Kalia VC. Phylogeny in aid of the present and novel microbial lineages: Diversity in *Bacillus*. *PLoS One* 2009; 4: e4438.
- [46] Pevzner PA, Sze SH. Combinatorial approaches to finding subtle signals in DNA sequences. In: Bourne PE, Gribskov M, Altman RB, Jensen N, Hope D, Lengauer T, Mitchell JC, Scheeff ED, Smith C, Strande S, Weissig H, eds. *Proceedings of the Eighth International Conference on Intelligent Systems for Molecular Biology*. AAAI Press, Menlo Park, CA., 2000; pp 269-278.
- [47] Stothard P. The Sequence Manipulation Suite: JavaScript programs for analyzing and formatting protein and DNA sequences. *Biotechniques* 2000; 28: 1102-1104.
- [48] van Baarlen P, van Belkum A, Summerbell RC, Crous PW, Thomma BPHJ. Molecular mechanisms of pathogenicity: how do pathogenic microorganisms develop cross-kingdom host jumps? *FEMS Microbiol Rev* 2007; 31: 239-277.
- [49] Vianna ME, Conrads G, Gomes BP, Horz HP. Identification and quantification of archaea in primary endodontic infections. *J Clin Microbiol* 2006; 44: 1274-1282.
- [50] Park S-Y, Kang H-O, Jang H-S, Lee J-K, Koo B-T, Yum DY. Identification of extracellular N-acylhomoserine lactone acylase from a *Streptomyces* sp. and its application to quorum quenching. *Appl Environ Microbiol* 2005; 71: 2632-2641.
- [51] Hewitt L, Kasche V, Lummer K, *et al.* Structure of a slow processing precursor penicillin acylase from *Escherichia coli* reveals the

- linker peptide blocking the active-site cleft. *J Mol Biol* 2000; 302: 887-898.
- [52] Miyazaki S, Sugawara H, Ikeo K, Gojobori T, Tateno Y. DDBJ in the stream of various biological data. *Nucleic Acids Res* 2004; 32: D31-D34.
- [53] Bai F, Han Y, Chen J, Zhang X-H. Disruption of quorum sensing in *Vibrio harveyi* by the AiiA protein of *Bacillus thuringiensis*. *Aquaculture* 2008; 274: 36-40.
- [54] Roche DM, Byers JT, Smith DS, Glansdorp FG, Spring DR, Welch M. Communications blackout? Do N-acylhomoserine-lactone-degrading enzymes have any role in quorum sensing? *Microbiology* 2004; 150: 2023-2028.
- [55] Leadbetter JR, Greenberg EP. Metabolism of acyl-homoserine lactone quorum-sensing signals by *Variovorax paradoxus*. *J Bacteriol* 2000; 182: 6921-6926.
- [56] Park SY, Lee SJ, Oh TK, Oh JW, Koo BT, Yum DY, Lee JK. AhlD, an N-acylhomoserine lactonase in *Arthrobacter* sp., and predicted homologues in other bacteria. *Microbiology* 2003; 149: 1541-1550.
- [57] Yang WW, Han JI, Leadbetter JR. Utilization of homoserine lactone as a sole source of carbon and energy by soil *Arthrobacter* and *Burkholderia* species. *Arch Microbiol* 2006; 185: 47-54.
- [58] Ochsner UA, Wilderman PJ, Vasil AI, Vasil ML. GeneChip expression analysis of the iron starvation response in *Pseudomonas aeruginosa*: Identification of novel pyoverdine biosynthesis genes. *Mol Microbiol* 2002; 45: 1277-1287.
- [59] Rasmussen TB, Givskov M. Quorum-sensing inhibitors as anti-pathogenic drugs. *Int J Med Microbiol* 2006; 296: 149-161.
- [60] Hentzer M, Wu H, Andersen JB, et al. Attenuation of *Pseudomonas aeruginosa* virulence by quorum sensing inhibitors. *EMBO J* 2003; 22: 3803-3815.
- [61] Rasmussen TB, Givskov M. Quorum sensing inhibitors: a bargain of effects. *Microbiology* 2006; 152: 895-904.

Received: December 04, 2010

Revised: December 28, 2010

Accepted: December 30, 2010

© Kalia et al.; Licensee Bentham Open.

This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.